



HAL
open science

Adjunct therapies to improve outcomes after botulinum toxin injection in children: A systematic review

L. Mathevon, Isabelle Bonan, J.-L. Barnais, F. Boyer, Mickaël Dinomais

► To cite this version:

L. Mathevon, Isabelle Bonan, J.-L. Barnais, F. Boyer, Mickaël Dinomais. Adjunct therapies to improve outcomes after botulinum toxin injection in children: A systematic review. *Annals of Physical and Rehabilitation Medicine*, Elsevier Masson, 2019, 62 (4), pp.283-290. 10.1016/j.rehab.2018.06.010 . hal-02535033

HAL Id: hal-02535033

<https://hal.univ-angers.fr/hal-02535033>

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial | 4.0 International License

Adjunct therapies to improve outcomes after botulinum toxin injection in children: a systematic review

L. Mathevon¹, I. Bonan², J-L. Barnais³, F. Boyer⁴, M. Dinomais⁵

¹ CHU de Grenoble, Department of Physical and Rehabilitation Medicine, France

² CHU de Rennes, Department of Physical and Rehabilitation Medicine, University of Rennes, France

³ CHU de Martinique, Department of Physical and Rehabilitation Medicine.

⁴ CHU de Reims, Department of Physical and Rehabilitation Medicine, University of Reims, France

⁵ CHU de Angers & Les Capucins Angers, Department of Physical and Rehabilitation Medicine, University of Angers, France

Corresponding author:

Laure MATHEVON

9 bis impasse de l'Aigas,

69160 Tassin la Demi-Lune

06.89.53.16.63; l.mathevon@laposte.net

Abstract

Background. Botulinum toxin (BTX) injection alone is not sufficient to treat spasticity in children, notably those with cerebral palsy; thus, there is an emerging trend for adjunct therapies to offer greater outcomes than BTX alone.

Objective. The aim of this systematic review was to evaluate the general effectiveness of adjunct therapies regardless of their nature in children with spasticity.

Methods. MEDLINE, Cochrane and Embase databases were searched from January 1980 to March 15, 2018 for reports of parallel-group trials (randomized controlled trials [RCTs] and non-RCTs) assessing adjunct therapies after BTX injection for treating spasticity in children. Two independent reviewers extracted data and assessed the risk of bias by using the PEDro scale for RCTs and Downs and Black scale (D&B) for non-RCTs.

Results. Overall, 20 articles involving 662 participants met the inclusion criteria. The average quality was good for the 16 RCTs (mean PEDro score 7.4 [SD 1.6]) and poor to moderate for the 4 non-RCTs (D&B score 9 to 17). Adjunct therapies consisted of casting/posture,

electrical stimulation, resistance training and rehabilitation programmes. Casting associated with BTX injection improved the range of passive and active motion and reduced spasticity better than did BTX alone (9 studies), with a follow-up of 1 year. Resistance training enhanced the quality and performance of muscles without increasing spasticity. Only 3 rehabilitation programmes were studied, with encouraging results for activities.

Conclusion. Lower-limb posture with casting in children has a high level of evidence, but the long-term efficacy of short-leg casting needs to be evaluated. A comparison between the different modalities of casting is missing, and studies specifically devoted to testing the different kinds of casting are needed. Moreover, the delay to casting after BTX injection is not clear. Data on electrical stimulation are not conclusive. Despite the small number of studies, resistance training could be an interesting adjunct therapy notably to avoid loss of strength after BTX injection. Rehabilitation programmes after BTX injection still need to be evaluated.

Key Words: Spasticity, Cerebral Palsy, Botulinum toxin, Casting, Rehabilitation programme

Introduction

As for adults, for children, spasticity is a common consequence of upper motor-neuron disorders due to spinal cord injury, traumatic brain injury, stroke, brain tumour and cerebral palsy (CP). CP, defined as a permanent motor disorder due to non-progressive damage to the developing brain, is a major cause of motor disability in children and accounts for most of the spasticity cases seen in children, in contrast to adults, in whom CP is not the major cause of spasticity. Among these CP children, approximately 80% to 90% have spastic forms (unilateral or bilateral) that result in abnormal motor function. In addition to a motor deficit cause, spasticity contributes to reduced motor ability¹⁻⁴ and is a cause of musculoskeletal deformities in children and *a fortiori* in children with CP.^{5,6}

Despite no cure for CP, motor impairment can be minimized with neuro-rehabilitation. An increasing number of studies in the last decade have highlighted multiple interventions in children with CP. A recent systematic review revealed what does and does not work, to help clinicians keep up to date and thus provide children and their families with effective, evidence-based interventions in motor rehabilitation^{7,8}.

Current rehabilitation interventions to treat motor impairments in CP are mainly based on techniques aimed at repeatedly stimulating the paretic limb and hence reducing spasticity^{9,10}. Protection of the musculoskeletal system also remains a major issue because it is required to preserve motor functions at the end of the growth period and prevent early ageing⁶ of the joints.

Botulinum toxin type A (BTX-A) is one of the most effective and safest treatments for focal spasticity in children with CP⁸ and more generally children with spasticity due to other causes. BTX reduces spasticity^{8,9} and maintains a favourable range of motion so as to prevent vicious joint patterns^{11,12}.

However, the level of evidence for the improvement of motor function with BTX is weak^{10,13}. One explanation is that BTX injection must be associated with a specific, personalised rehabilitation programme (adjunct therapies). A large heterogeneity of practices concern adjunct therapies after BTX injections to optimise the results. Although BTX treatment is highly recommended to treat focal spasticity (grade A level of evidence⁸) in children with CP (and by extrapolation to treat focal spasticity in children), no clear data are available to provide recommendations for the optimal adjunct therapies after such treatment.

The aim of this study was to evaluate, via a systematic literature review, the general effectiveness of all types of adjunct therapies used after BTX injection on outcomes related to impaired body function and structure limited to the activity and participation of children with spasticity.

Methods

Study eligibility criteria

Three authors (LM, JLB and MD) performed a systematic review of the literature in accordance with PRISMA guidelines (www.prisma-statement.org). We included any English reports of adjunct therapies after BTX injection to treat spasticity in children. Only parallel clinical studies (i.e., comparing least 2 groups [adjunct vs no adjunct therapies]) were included. Articles concerning adjunct therapies after phenolisation were excluded because since 2009, phenolisation is no longer recommended for treating spasticity in children¹⁴. Literature reviews were not analysed.

MEDLINE, Cochrane, PEDro and Reeduc databases were searched from 1980 to March 15, 2018 with the key words: “botulinum toxin” AND “children” (OR “child”, “kids”, “childhood”, “infant”, “adolescent”) AND “spasticity”. We deliberately chose not to use the term “cerebral palsy” because spasticity is also present in children after stroke, traumatic brain injury etc. (see Introduction). Related articles and links were also searched. Additional articles were identified by a manual search of references in key articles retrieved.

Quality analysis

The quality of RCTs was analysed by the specific PEDro scale¹⁵. The score ranged from 0 to 11. A score of 0 to 4 was considered poor quality, 5 or 6 moderate, 7 or 8 good, and ≥ 9 excellent. The quality of non-RCTs was analysed by the specific Downs and Black scale (D&B)¹⁶, with scores ranging from 0 to 27 (high quality). The following cut-points have been suggested to classify studies by quality: excellent (26–27), good (20–25), fair (15–19) and poor (0–18)¹⁷. Quality was analysed independently by both LM and JLB, with any disagreements resolved after discussion with a third evaluator (MD). To provide the reader with a clear overview of our findings, we assigned each adjunct therapy to 1 of 3 categories: “do it”; “probably do it”, or “don’t do it”, inspired by Novak et al.⁸

Results (Table 1)

Among 519 articles published from January 1980 to March 15, 2018 from the 4 data sources (Fig. 1), after reading the titles and abstracts, 39 articles were initially selected. After reading the full text, 19 articles were excluded; 20 articles were finally deemed relevant. At the time of this work, no previous systematic review had focused on adjunct therapies after BTX injection in children. All 20 articles had been published between 2001 and March 2018, 9 after 2010; 16 were RCTs and 4 prospective interventional studies.

Quality of articles

The quality analysis is presented in Table 1. The mean PEDro quality score for the 16 RCTs was 7.4 (SD 1.6; range 4–9), for good quality. The D&B quality score for the 4 non-RCTs ranged from 9 to 17, for poor to moderate quality. The total number of participants included was 662, with a large sample size (> 50) in 3 studies^{18–20}. The number of subjects needed for statistical power was not calculated in the articles retained, apart from 2 studies with 50% and 60% of the calculated sample size recruited^{21,22}.

CP was exclusively investigated in all studies; we found no studies related to other aetiologies of spasticity in children. The motor deficit was mainly mixed unilateral and bilateral in 9 studies, exclusively unilateral in 6 studies, and exclusively bilateral in 1 study. The functional level of participants was good. Upper-limb function was investigated in 5 studies, with a Manual Ability Classification System level reported as 1 to 3 in 2 studies^{22,23}. The other studies investigated lower-limb function, with a Gross Motor Function Classification System (GMFCS) level of at least 3 and at least 2 in 7 studies.

Every study fully described the experimental protocol. Most studies compared adjunct therapy associated with BTX injection versus BTX alone. In 2 studies^{21,24}, adjunct therapy was investigated alone in comparison with BTX ± adjunct therapy. In terms of the components of the World Health Organization's International Classification of Functioning, Disability and

Health (ICF, WHO 2001), all studies focused on outcomes related to body structure and function, such as range of motion, spasticity, and muscle strength. For activity, the Gross Motor Function Measure (GMFM) was used in 7/15 studies that investigated lower-limb function. Upper-limb function was investigated with many scales with good metrological properties (Table 1). Only one RCT focused on quality of life with the Child Health Questionnaire (CHQ)²⁵.

Meta-analysis is generally considered a more powerful estimate of the true effect size than a single study. However, meta-analysis was precluded in our study because of the heterogeneity of study designs regarding the different kinds of adjunct therapies (posture, serial casting, casting or splints etc.), diversity of participants (different ages, different inclusion criteria, localisation of spasticity etc.) and number and variability of outcomes studied (impairment [range of motion, spasticity etc.] and activities [Assisting Hand Assessment, GMFM etc.]).

Synthesis of the literature

The different adjunct therapy protocols proposed in the available literature were classified as relating to posture (n=10 articles), electrical stimulation (n=4) and rehabilitation procedures (n=6).

Joint Posture Procedure

Different joint posture procedures were reported. Six articles adopted serial casting protocols and 3 evaluated the effects of fixed casting. Potential confounding factors such as day or night posture and duration of casting were not systematically mentioned and therefore could not be taken into account. Also, because length of follow-up never exceeded 12 months, long-term effects could not be evaluated.

Serial casting (n=6)

This consisted of applying 3 progressive casts: 5 studies used a short-leg cast^{18,20,24,26,27} and the other a long-leg cast²⁵. The timing of serial cast application ranged from immediate²⁶ to 4 weeks²⁷, 1 week¹⁸ or 3 weeks^{20,25} or 1 to 3 weeks²⁴ after BTX injection in the triceps surae^{18,20,24,26,27}. Casts were applied continuously except in one study¹⁸ in which casts were applied for 72 hr/week. Only 1 study²⁴ compared BTX injection with casting versus casting alone. The quality of the studies was good for the 4 RCTs, with an average PEDro score of 8.3^{18,24,27,25}, and moderate for the 2 non-RCTs^{20,26}, with an average D&B score of 16.5^{20,26}. The total number of participants was 280. All 6 studies evaluated body structure and function outcomes. Only one RCT focused on quality of life with the CHQ.

Whatever the protocol of serial casting, the improved passive range of motion and the reduced spasticity (Modified Ashworth Scale [MAS] and Tardieu Scale [TS]) were better when associating casting with BTX than BTX alone, even when the casts were applied sequentially¹⁸. Kay et al.²⁴ showed that the reduced spasticity (MAS) was maintained longer (up to 1 year) with casting without than with BTX injection (3 months). In this study, serial casting was applied until $\geq 5^\circ$ of dorsiflexion was reached with the knee extended, then fixed casting was applied during follow-up. This protocol could be considered a prolonged mixed casting regimen, so one cannot discriminate specific serial casting effects from those of other interventions (prolonged fixed casting or BTX alone). No conclusions could be drawn from this result.

Concerning activity measurements related to gross motor function and/or functional mobility, very few improvements were reported. In the short-leg cast protocol, GMFM scores were initially unchanged^{24,26} but started to increase after the first 3 months of treatment, with no significant difference between the 2 groups in 1 study²⁴. In the long-leg cast protocol, GMFM and CHQ scores were markedly elevated in both groups after 6 weeks²⁵.

Some adverse effects from serial casting were noted: pain and atrophy (measured from calf and thigh perimeters) were higher **versus** BTX injection alone^{27,25}. The casts also **seemed** more difficult to bear immediately after BTX injection than 4 weeks later, with no difference when evaluating body structure and function outcomes²⁷.

Casting (n=4)

Continuous fixed casting consisted **of applying** one short-leg cast immediately after BTX injection in the triceps surae for 2 weeks²⁸ or 3 weeks^{21,29} **and** for single²⁹ or multiple sessions of treatment^{21,28} with a long follow-up (at least 8 months). All treatment groups had physiotherapy several times a week **after** BTX injection. The quality of the studies was good for 2 **studies** (PEDro scores 9 and 7)^{22,28} and poor (PEDro score 5) for the third **RCT**²⁹, for a total of 69 patients.

Concerning outcomes representing the ICF component of body functions and structures, casting associated with BTX injection and even casting alone improved passive range of motion and ankle motor selectivity, reduced spasticity (MAS, TS)^{21,28} and improved gait parameters as measured by the GMFM²⁸ or Observational Gait Scale²⁸ and even improved the spatiotemporal parameters and ankle kinematics **on** quantified gait analysis²¹. These results were maintained in the relative long term with 2 or 3 sessions of treatment (between 8 months and 1 year). **Of note, the** use of BTX alone **conferred** no improvement in terms of impairment and gait parameters²¹. No adverse effects were reported.

The use of a night splint **after** BTX injection in the upper limb was evaluated in a single RCT **with** poor quality (PEDro score 4)³⁰ and surprisingly, suggested that upper-limb function (**Quality of Upper Extremity Skills Test** score) was significantly better **than with** BTX alone after 6 months of regular use ($p < 10^{-4}$).

Electrical stimulation (n=4)

Two kinds of **protocols** **were found**.

The first^{31,32} (n=2) used electrical stimulation in the gastrocnemius muscles after BTX injection in the triceps surae **versus** BTX alone. Different modalities of stimulation exist in terms of type of current (continuous or rectangular biphasic, high or low frequency), duration (15 to 30 **min**) and frequency of application (1-6 times/day for 3 days to 6 weeks). The quality of these 2 RCTs was good (PEDro score 8 for both) but with **only 35 total participants**. Electrical stimulation **did not** provide any additional effects **as** compared with BTX alone in terms of **reduced** spasticity or **improved** gait (Physician Rating Scale, 3D Instrumented Gait analysis).

The second protocol^{33,34} (n=2) consisted **of** functional electrical stimulation in antagonist muscles after BTX injection in the wrist and finger flexors³³ and in the triceps surae³⁴ associated with a physical therapy programme. The quality of these 2 studies was poor (PEDro **score 5** and D&B **score 9, respectively**) and no conclusions can be drawn.

Multimodal Rehabilitation Procedures

Resistance training (n=3)

This **training** consisted of a global progressive programme tailored to each individual for strengthening the spastic and antagonist muscles in the lower^{35,36} or upper limb²³ twice or 3 times/week for 8 to 12 weeks. These programmes were compared to physiotherapy without resistance training (gait, balance, stretching) with good methodological quality^{35,36} (PEDro **score 9**) or nothing **after** BTX injection in the upper limb, with moderate quality (PEDro **score 6**)²³. The total number of participants was 39. Resistance training improved muscle strength in injected muscles³⁵ without increasing spasticity. Williams et al. attributed the decrease in muscle atrophy **after** BTX observed on MRI to the resistance training³⁵. These RCTs showed no impact on body functions, structures or activity (assisting hand assessment²³, biomechanical force plate³⁶, gait parameters with GMFM and instrumented gait analysis³⁶), but personalised goals were achieved (**Goal Attainment Scaling**)³⁵.

Rehabilitation programmes (n=3)

Surprisingly, only 3 studies evaluated a rehabilitation programme after BTX injection: Modified Constraint Induced Therapy³⁷ (mCIMT) (n=29, D&B score 17) and Bimanual Task Oriented Therapy (BTOT)²² (n=35, PEDro score 8) for the upper limb and a poorly documented global rehabilitation programme for 2 hr/day¹⁹ (n=244, PEDro score 8) for the lower limb.

“Rehabilitation” associated with the BTX injection conferred more improvement in the functional scales than with BTX alone: GMFM¹⁹, Upper Limb Physician's Rating Scale, and How Often scale and How Well scale in the revised Paediatric Motor Activity Log³⁷. Of note, 6 months after the treatment, functional grip strength was significantly better with BTOT alone than with BTOT associated with BTX injection²².

Discussion

BTX injection alone is not sufficient to reach functional goals during a child's growth⁸. Surprisingly, although the number is increasing, few studies have been published on adjunct therapies after BTX injection during the last decade³⁸. Findings from this systematic review of the effectiveness of adjunct therapies are based on 20 studies involving 662 children with CP, essentially concerning posture, with only 6 studies investigating specific rehabilitation after BTX injection (3 of the upper limb). Nevertheless, the quality of these studies is good.

However, the number of participants needed for statistical power was not calculated or reported in any study included in our review, which reduces the strength of any conclusions that can be drawn from the results. Concerning outcome assessment in the studies included, few studies focused on outcomes related to activities, with most reporting outcomes related to only body structure and function. This finding is similar to the previous conclusions of Tustin et al.³⁹ and Blackmore et al.³⁸

This systematic review confirms that BTX injections alone are not sufficient to improve outcomes⁸, and adjunct therapies are clearly necessary: BTX injections alone cannot be recommended for children.

During growth, lower-limb posture with casting is recommended with a high level of evidence, although only the short-term efficacy of short-leg casting has been evaluated. To date, no study has compared the different modalities of casting, and further studies specifically devoted to testing these different kinds of casting are needed. No clear recommendation can be made concerning the delay after BTX injection before applying casting. Some authors have proposed that BTX begins to take effect 1 week after injection, but **we found** no evidence for this. Serial casting has not demonstrated greater efficacy than a single cast that should be worn for at least 2 weeks. In a **recently** published work⁴⁰ (not **included** in our analysis because the paper was published after our endpoint), the authors did not show any significant differences between serial or single casting after BTX injection in the triceps surae, and **the** magnitude of improvements **was** similar between single and serial casting procedures. If confirmed in **a** large sample, applying single casting **may be preferred because of** the greater convenience for both clinicians and children.

Moreover, adverse effects such as pain, muscle atrophy and cutaneous lesions **were noted** in serial casting rather than unique casting **studies**. **Although clear** evidence exists for the lower limb, the sparse number of studies for posture **therapy** in the upper limb **after** BTX **injection** in children with CP provides only a moderate level of evidence. However, we can extrapolate from the available literature concerning lower-limb spasticity treatment that posture **therapy** should also be recommended after BTX injection to treat spasticity in upper limbs.

Personalised resistance training involving both spastic and antagonist muscles and depending on individualised motor control helped reach personalised goals and treat spasticity. Moreover, adding resistance training **seems** able to control muscle atrophy **after** BTX injection. However,

the sample size of these studies was very limited. Further studies with a larger sample size are needed. Moreover, the timing of a **resistance training** programme is **unknown because such training** pre- or post-BTX injection probably **has** different goals. **The training** before injection should help protect muscles **against** atrophy, whereas **that after** BTX aims to work on motor control in a personalised way³⁵.

In the upper limb, mCIMT and BTOT **had** positive results in terms of function with or without BTX injection. Further studies should be conducted with the use of BTX injection pre-mCIMT or BTOT only if spasticity is a major impairment regarding upper limb function.

Because of few studies for goal-directed rehabilitation programmes **after** BTX injection, we have only a moderate level of evidence for adjunct therapies. However the available literature^{7,8} regarding motor rehabilitation for **children with** CP after BTX injection considers such adjunct goal-directed approaches promising.

Finally, the results obtained with electrical stimulation are not encouraging, although the number of participants in these studies was very low. Electrical stimulation as adjunct therapy after BTX injection in children **seems** inefficient.

Conclusions (see Table 2)

Adjunct therapies such as posture (serial casting, casting or splints), goal-directed rehabilitation programmes and strength training may improve spasticity outcomes **in children** when used after BTX injection. This **therapy** has been clearly demonstrated for **treating** lower-limb spasticity. The level of evidence is low **for the effectiveness of** BTX injection without adjunct therapy **for** impairment (**range of motion**, spasticity) and activities (notably gait parameters). **However**, a high level of evidence **suggests** that posture **therapy** after BTX **injection** **improves range of motion**, spasticity and gait parameters. Further research is needed to test the best procedure for posture (serial casting, casting or splints) and to determine the optimal time window for applying these adjunct therapies. **The** level of evidence **is moderate** for small to moderate

improvements in impairment (range of motion, spasticity) and activities (gross motor function) for resistance training and rehabilitation programmes. Finally, the level of evidence is low for the effectiveness of electrical stimulation. To give a clear overview of our conclusions, for each adjunct therapy listed in Table 2, we have provided a short comment and assigned it to one of three categories: “do it”; “probably do it”, or “don’t do it”.

ICF domains of activities, participation and quality of life need to be considered in future studies to establish the clinical relevance of adjunct therapies in children. Finally, no study was conducted in children with spasticity due to other neurological conditions. Hence, the conclusions are strictly applicable to only children with CP. Further studies are needed to test adjunct therapies after BTX injection in children with other causes of spasticity.

Conflict of interest: None declared.

Titles and legends

Figure. Flow of studies in the review.

Table 1. Synthesis of the 20 selected studies: type of study, population, methods and results. Results are presented as mean \pm SD or (min-max) Group 1 vs Group 2 (\pm Group 3) for pre- and post-treatment values at the end of follow-up. *P < 0.05, **P \leq 0.001 compared to pretreatment values

When the mean change between pre- and post-treatment was calculated for each group, it is presented with SD values † p < 0.05 (difference between groups)

RCT, randomized controlled trial; BTX, botulinum toxin type A; MAS, Modified Ashworth Scale; TS, Tardieu scale; PROM, Passive Range of Motion in degrees; R1, spastic catch in Tardieu’s scale in degrees; OGS: Observational Gait Scale; PGA, Physician Global Assessment; QUEST, Quality of Upper Extremity Skills Test; MACS, Manual Ability Classification System; GMFM, Gross Motor Function Measure; GMFCS, Gross Motor Function Classification System; GAS, Goal Attainment Scaling
Speed gait in km.h⁻¹ or cm.min⁻¹ or m.s⁻¹; stride length in cm; maximal torque in 0.1 Nm/kg; functional strength in g

Table 2. Synthesis of level of evidence.

References


1. Chaleat-Valayer E, Bard-Pondarre R, Bernard JC, et al. Upper limb and hand patterns in cerebral palsy: Reliability of two new classifications. *Eur J Paediatr Neurol EJPJN Off J*

- Eur Paediatr Neurol Soc.* 2017;21(5):754-762.
2. Palisano RJ, Hanna SE, Rosenbaum PL, Tieman B. Probability of walking, wheeled mobility, and assisted mobility in children and adolescents with cerebral palsy. *Dev Med Child Neurol.* 2010;52(1):66-71.
 3. Beckung E, Hagberg G, Uldall P, Cans C, Surveillance of Cerebral Palsy in Europe. Probability of walking in children with cerebral palsy in Europe. *Pediatrics.* 2008;121(1):e187-e192.
 4. Beckung E, Hagberg G. Neuroimpairments, activity limitations, and participation restrictions in children with cerebral palsy. *Dev Med Child Neurol.* 2002;44(5):309-316.
 5. Rethlefsen SA, Blumstein G, Kay RM, Dorey F, Wren TAL. Prevalence of specific gait abnormalities in children with cerebral palsy revisited: influence of age, prior surgery, and Gross Motor Function Classification System level. *Dev Med Child Neurol.* 2017;59(1):79-88.
 6. Horstmann HM, Hosalkar H, Keenan MA. Orthopaedic issues in the musculoskeletal care of adults with cerebral palsy. *Dev Med Child Neurol.* 2009;51 Suppl 4:99-105.
 7. Ward R, Reynolds JE, Bear N, Elliott C, Valentine J. What is the evidence for managing tone in young children with, or at risk of developing, cerebral palsy: a systematic review. *Disabil Rehabil.* 2017;39(7):619-630.
 8. Novak I, McIntyre S, Morgan C, et al. A systematic review of interventions for children with cerebral palsy: state of the evidence. *Dev Med Child Neurol.* 2013;55(10):885-910.
 9. Strobl W, Theologis T, Brunner R, et al. Best Clinical Practice in Botulinum Toxin Treatment for Children with Cerebral Palsy. *Toxins.* 2015;7(5):1629-1648.
 10. Fehlings D, Novak I, Berweck S, et al. Botulinum toxin assessment, intervention and follow-up for paediatric upper limb hypertonicity: international consensus statement. *Eur J Neurol.* 2010;17 Suppl 2:38-56.
 11. Chaléat-Valayer E, Parratte B, Colin C, et al. A French observational study of botulinum toxin use in the management of children with cerebral palsy: BOTULOSCOPE. *Eur J Paediatr Neurol EJPN Off J Eur Paediatr Neurol Soc.* 2011;15(5):439-448.
 12. Barrett RS. What are the long-term consequences of botulinum toxin injections in spastic cerebral palsy? *Dev Med Child Neurol.* 2011;53(6):485.
 13. Hoare BJ, Wallen MA, Imms C, Villanueva E, Rawicki HB, Carey L. Botulinum toxin A as an adjunct to treatment in the management of the upper limb in children with spastic cerebral palsy (UPDATE). *Cochrane Database Syst Rev.* 2010
 14. Simon O, Yelnik AP. Managing spasticity with drugs. *Eur J Phys Rehabil Med.* 2010;46(3):401-410.
 15. Moseley AM, Herbert RD, Sherrington C, Maher CG. Evidence for physiotherapy practice: a survey of the Physiotherapy Evidence Database (PEDro). *Aust J Physiother.* 2002;48(1):43-49.
 16. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health.* 1998;52(6):377-384.
 17. Hooper P, Jutai JW, Strong G, Russell-Minda E. Age-related macular degeneration and low-vision rehabilitation: a systematic review. *Can J Ophthalmol J Can Ophthalmol.* 2008;43(2):180-187.
 18. Dursun N, Gokbel T, Akarsu M, Dursun E. Randomized Controlled Trial on Effectiveness of Intermittent Serial Casting on Spastic Equinus Foot in Children with Cerebral Palsy After Botulinum Toxin-A Treatment. *Am J Phys Med Rehabil.* 2017;96(4):221-225.
 19. Jianjun L, Shurong J, Weihong W, Yan Z, Fanyong Z, Nanling L. Botulinum toxin-A

- with and without rehabilitation for the treatment of spastic cerebral palsy. *J Int Med Res.* 2013;41(3):636-641.
20. Lee SJ, Sung IY, Jang DH, Yi JH, Lee JH, Ryu JS. The effect and complication of botulinum toxin type a injection with serial casting for the treatment of spastic equinus foot. *Ann Rehabil Med.* 2011;35(3):344-353.
 21. Ackman JD, Russman BS, Thomas SS, et al. Comparing botulinum toxin A with casting for treatment of dynamic equinus in children with cerebral palsy. *Dev Med Child Neurol.* 2005;47(9):620-627.
 22. Speth L, Janssen-Potten Y, Rameckers E, et al. Effects of botulinum toxin A and/or bimanual task-oriented therapy on upper extremity activities in unilateral Cerebral Palsy: a clinical trial. *BMC Neurol.* 2015;15:143.
 23. Elvrum A-KG, Brændvik SM, Sæther R, Lamvik T, Vereijken B, Roeleveld K. Effectiveness of resistance training in combination with botulinum toxin-A on hand and arm use in children with cerebral palsy: a pre-post intervention study. *BMC Pediatr.* 2012;12:91.
 24. Kay RM, Rethlefsen SA, Fern-Buneo A, Wren TAL, Skaggs DL. Botulinum toxin as an adjunct to serial casting treatment in children with cerebral palsy. *J Bone Joint Surg Am.* 2004;86-A(11):2377-2384.
 25. Dai AI, Demiryürek AT. Serial Casting as an Adjunct to Botulinum Toxin Type A Treatment in Children With Cerebral Palsy and Spastic Paraparesis With Scissoring of the Lower Extremities. *J Child Neurol.* 2017;32(7):671-675.
 26. Park ES, Rha D-W, Yoo JK, Kim SM, Chang WH, Song SH. Short-term effects of combined serial casting and botulinum toxin injection for spastic equinus in ambulatory children with cerebral palsy. *Yonsei Med J.* 2010;51(4):579-584.
 27. Newman CJ, Kennedy A, Walsh M, O'Brien T, Lynch B, Hensey O. A pilot study of delayed versus immediate serial casting after botulinum toxin injection for partially reducible spastic equinus. *J Pediatr Orthop.* 2007;27(8):882-885.
 28. Hayek S, Gershon A, Wientroub S, Yizhar Z. The effect of injections of botulinum toxin type A combined with casting on the equinus gait of children with cerebral palsy. *J Bone Joint Surg Br.* 2010;92(8):1152-1159.
 29. Bottos M, Benedetti MG, Salucci P, Gasparroni V, Giannini S. Botulinum toxin with and without casting in ambulant children with spastic diplegia: a clinical and functional assessment. *Dev Med Child Neurol.* 2003;45(11):758-762.
 30. Kanellopoulos AD, Mavrogenis AF, Mitsiokapa EA, et al. Long lasting benefits following the combination of static night upper extremity splinting with botulinum toxin A injections in cerebral palsy children. *Eur J Phys Rehabil Med.* 2009;45(4):501-506.
 31. Rha D, Yang EJ, Chung HI, Kim HB, Park C, Park ES. Is electrical stimulation beneficial for improving the paralytic effect of botulinum toxin type A in children with spastic diplegic cerebral palsy? *Yonsei Med J.* 2008;49(4):545-552.
 32. Detrembleur C, Lejeune TM, Renders A, Van den Bergh PYK. Botulinum toxin and short-term electrical stimulation in the treatment of equinus in cerebral palsy. *Mov Disord Off J Mov Disord Soc.* 2002;17(1):162-169.
 33. Pieber K, Herceg M, Wick F, Grim-Stieger M, Bernert G, Paternostro-Sluga T. Functional electrical stimulation combined with botulinum toxin type A to improve hand function in children with spastic hemiparesis - a pilot study. *Wien Klin Wochenschr.* 2011;123(3-4):100-105.
 34. Seifart A, Unger M, Burger M. Functional electrical stimulation to lower limb muscles after botox in children with cerebral palsy. *Pediatr Phys Ther Off Publ Sect Pediatr Am Phys Ther Assoc.* 2010;22(2):199-206.
 35. Williams SA, Elliott C, Valentine J, Gubbay A, Shipman P, Reid S. Combining strength training and botulinum neurotoxin intervention in children with cerebral palsy: the


- impact on muscle morphology and strength. *Disabil Rehabil.* 2013;35(7):596-605.
36. Bandholm T, Jensen BR, Nielsen LM, et al. Neurorehabilitation with versus without resistance training after botulinum toxin treatment in children with cerebral palsy: a randomized pilot study. *NeuroRehabilitation.* 2012;30(4):277-286.
37. Park ES, Rha D-W, Lee JD, Yoo JK, Chang WH. The short-term effects of combined modified constraint-induced movement therapy and botulinum toxin injection for children with spastic hemiplegic cerebral palsy. *Neuropediatrics.* 2009;40(6):269-274.
38. Blackmore AM, Boettcher-Hunt E, Jordan M, Chan MDY. A systematic review of the effects of casting on equinus in children with cerebral palsy: an evidence report of the AACPDM. *Dev Med Child Neurol.* 2007;49(10):781-790.
39. Tustin K, Patel A. A Critical Evaluation of the Updated Evidence for Casting for Equinus Deformity in Children with Cerebral Palsy. *Physiother Res Int J Res Clin Phys Ther.* 2017;22(1).
40. Kelly B, MacKay-Lyons M, Berryman S, Hyndman J, Wood E. Casting Protocols Following BoNT-A Injections to Treat Spastic Hypertonia of the Triceps Surae in Children with Cerebral Palsy and Equinus Gait: A Randomized Controlled Trial. *Phys Occup Ther Pediatr.* May 2018:1-17.

519 total articles identified from
electronic databases
Medline, Cochrane and Embase
databases



480 articles excluded based on
titles and abstracts

39 full text articles assessed for
eligibility



19 full text articles excluded :
*10 articles didn't have a
comparison group
*4 study protocols
*2 systematic reviews
*3 articles out of topic

20 original articles included in
review

Authors	Design	Adjunct therapy	Therapy protocol	Population	Assessment of Deficiencies	Activity Outcomes	Quality of life	Follow-up	Results	PEdro	D&B
Dai et al. ²⁸ 2017	RCT	Serial Casting	3 progressive long leg castings Lower extremities placed in abduction First one 3 weeks after BTX-A injection Each for 1 week Group 1 : BTX + casting group Group 2 : BTX alone	70 children Cerebral Palsy Bilateral Scissoring of the lower extremities Mean age : 3.3 years	Spasticity: MAS	GMFM-66	Child Health Questionnaire (CHQ)	3 months	Pretreatment : MAS : 3.7±5.2 vs 3.8±3.4 GMFM-66 : 41.2±7.6 vs 42.3±2.5 CHQ : 42.7±2.9 vs 44.3±5.4 Posttreatment : MAS : 1.9±1.6 vs 2.8±6.3 GMFM-66 : 77.4±6.8 vs 64.6±3.5** CHQ : 76±6.7 vs 64.6±3.5** Mean change group 1 vs group 2 : MAS : 1.8 vs 1.0 GMFM-66 : 36.2 vs 22.3 CHQ : 33.3 vs 20.3	7	
Dursun et al. ¹⁸ 2017	RCT	Serial casting (intermittent)	3 progressive short leg castings First one 1 week after BTX-A injection Each for 72 hours Group 1 : BTX-A + casting group Group 2 : BTX alone	51 children Cerebral Palsy Unilateral / Bilateral GMFCS 1,2,3 Mean Age: 7.2 years	Spasticity: MAS, TS PROM	Gait: OGS Physician Global Assessment (PGA)		3 months	Pretreatment : MAS : 4.0 vs 4.0 PROM : 79.8±9.2 vs 79.4±9.0 OGS : 7.5±2.9 vs 8.5±2.7 PGA : N/A Posttreatment : MAS : 2.4±0.8 vs 3.1±0.9* PROM : 93±11.5 vs 83.5±12.7* OGS : 10.5±3.1 vs 9.5±3.0 PGA : 2.3±0.9 vs 1.3±0.7** Mean change group 1 vs group 2 :	8	
Lee et al. ²⁰ 2011	Prospective, Non RCT	Serial Casting	3 progressive consecutive short leg castings First one 3 weeks after BTX-A injection Each for 1 week Group 1 : BTX-A + casting group Group 2 : BTX alone	86 children Cerebral Palsy Unilateral / Bilateral GMFCS 1,2,3 Mean age: 5 years	Spasticity: MAS, TS PROM	Gait: Physician Rating Scale (PRS)		3 months	Pretreatment : MAS : 1.7±0.3 vs 1.5±0.4 PRM : -0.5±4.6 vs 0.7±5.2 PRS : 7.9±3.5 vs 8.2±3.4 Posttreatment : MAS : 1.2±0.3* vs 1.1±0.2 PRM : 9.0±3.4** vs 5.8±3.7* PRS : 11.3±2.4** vs 10.3±3.5* Mean change group 1 vs group 2 : MAS : 0.5 vs 0.4 PRM : 9.5 vs 5.1 PRS : 3.4 vs 2.1	17	
Park et al. ²⁹ 2010	Prospective, Non RCT	Serial Casting	3 progressive consecutive short leg castings First one immediately after BTX-A injection Each for 1 week Group 1 : BTX + casting group Group 2 : BTX alone	38 children Cerebral palsy Unilateral / Bilateral GMFCS 1,2,3 Mean age: 4.7 years	Spasticity: MAS, TS PROM	GMFM-66 (Dimension D and E)		1 month	Pretreatment : MAS : 2.9±0.8 vs 3.2±0.7 PROM : -1.2±1.5 vs 2.2±4.6 GMFM (D) : 75.3±29.6 vs 72.4±23.9 Posttreatment : MAS : 2.0±0.3* vs 2.3±0.7* PROM : 9.8±9.5* vs 7.2±6.30* GMFM (D) : 79.5±26.60* vs 73.5±19.3 Mean change group 1 vs group 2 : MAS : -1.0±0.9 vs -1.0±0.7 PROM : 11.0±9.8 vs 5.0 ± 6.1 † GMFM (D) : 4.2±5.7 vs 1.1±9.1	16	
Newman et al. ³⁰ 2007	RCT	Serial casting immediately or delayed	3 progressive consecutive short leg castings First one: 6 children immediately after BTX-A injection and 6 children 4 weeks later Each for 1 week Group 1 : BTX + immediate serial casting Group 2 : BTX + delayed serial casting	12 children Cerebral Palsy Unilateral/bilateral GMFCS 1,2 Mean age: 5.3 years	Spasticity: TS	Gait: Observational Gait Scale (OGS)		6 months	Pretreatment : R1 : -24.3±12.0 vs -25.7±5.3 PROM : -6.0±7.7 vs 3.3±4.0 OGS : 9.4±2.1 vs 11.2±2.7 Posttreatment : R1 : -21.1±11.2 vs -7.1±7.6** PROM : 0.0±4.8 vs 2.6±7.1* OGS : 10.3±5.3 vs 13.4±3.3** Mean change group 1 vs group 2 : R1 : 3.1±11.0 vs 18.6±10.4† PROM : 6.0±9.2 vs 6.4±6.0 OGS : 0.9±5.3 vs 2.2±2.8	8	
Kay et al. ²⁷ 2004	RCT	Serial Casting	Progressive consecutive short leg casting changed every two weeks until 25° of dorsiflexion was reached with knee extended Group 1 : Casting + BTX Group 2 : Casting only	23 children Cerebral Palsy Unilateral / Bilateral GMFCS 1,2,3 Mean age: 7.1 years	Spasticity: MAS PROM Computerized gait analysis: peak dorsiflexion (PD) swing	GMFM (C,D,E)		1 year	Pretreatment : MAS : 2.6±1.2 vs 2.6±1.1 PROM : -6.4±8.3 vs -3.7±8.7 PD swing : -12.3±1.1 vs -16.9±15.1 GMFM : 75.8±20.1 vs 66.4±23.1 Mean change group 1 vs group 2 : MAS : -0.9±1.0* vs -1.1±1.2* PROM : 18.4±11.7* vs 13.9±11.8** PD swing : 12.5±9.3* vs 15.1±11.8** GMFM : 2.5±7.5 vs -1.3±5.1	9	
Hayek et al. ⁴¹ 2010	RCT	Casting	Casts were applied on the day of the first injection and retained for 2 weeks.	20 children Cerebral Palsy Unilateral / Bilateral GMFCS : 1,2,3 Mean age: 13.5 years	Spasticity: TS PROM Selective motor control of the ankle	Gait analysis: Spatio-temporal parameters OGS GMFM-66		8 months	Pretreatment : R1 : -20.0±13.9 vs -21±12.6 PROM : 3.5±12.5 vs 8.7±7.5 OGS : N/A Gait Speed : 0.6±0.1 vs 0.6±0.1 GMFM : 54.0±15.4 vs 52.2±15.6 Mean change group 1 vs group 2 : R1 : -12.3±3.4** vs -13.3±3.8** PROM : 9.8±9** vs 11.9±7.5 † OGS : 12.6±1.4** vs 10.5±1.5** † Speed : 0.9±0.3** vs 0.9±0.3** GMFM : 64.3±3.7** vs 55.6±4.8**	7	
Ackman et al. ²⁶ 2005	Multicentre, RCT	Casting	Casts were applied on the day of the first injection and retained for 3 weeks. 3 cycles of treatment Group 1 : BTX alone Group 2 : Placebo injection + casting Group 3 : BTX + casting	39 children Cerebral Palsy Unilateral / bilateral GMFCS 1,2 Mean age: 6 years	Spasticity: MAS, TS PROM, AROM Ankle dorsiflexion and plantar flexion strength Ankle power generation Ankle kinematics	Gait analysis: Spatio-temporal parameters		1 year	N/A (results presented in diagrams) Group 1 : no significant change Group 2 and group 3 : significant improvements in spasticity, PROM and dorsiflexion strength and ankle kinematics.	9	
Bottos et al. ⁴² 2003	RCT	Casting	Casts applied on the day of the first injection and retained for 3 weeks. Group 1 : BTX + casting group Group 2 : BTX alone	10 children Cerebral Palsy Bilateral GMFCS 1,2,3 Mean age: 6.4 years	Spasticity: MAS	GMFM Computerized gait analysis		1 year	Mean change group 1 vs group 2 at 4 months : MAS : N/A (diagrams) GMFM : N/A (diagrams) → Both improved significantly in group 1 Stride length : 10.5 (N/A) vs 5.5 (N/A) † Speed walking : (cm.s ⁻¹) 5.3(N/A) vs 0.6 †	5	
Kanellopoulos et al. ⁴⁴ 2009	RCT	Night splint	Thermoplastic night splint applied 6 months after 2 BTX injections in the upper limb Group 1 : BTX + night splint Group 2 : BTX alone	20 children Cerebral Palsy Unilateral MACS ? Mean age: 7 years		Quality of Upper Extremity Skills Test (QUEST)		6 months	Mean change group 1 vs group 2 (N) : QUEST : 15.9 vs 4.2 †	4	
Pieber et al. ⁴³ 2011	RCT	Functional electrical stimulation (FES)	Stimulation of wrist and hand extensor muscles started 5 to 7 days after the injection Twice a day, 15 min for 3 months Biphasic rectangular current with a frequency of 30 Hz, 0.2 ms pulse width, 2-5 s on time Group 1 : BTX + FES Group 2 : BTX alone	6 girls Cerebral Palsy Unilateral MACS ? Mean age: 11.7 years	Spasticity: MAS, TS PROM AROM BMRC scale	Movement ABC checklist		6 months	Mean change Group 1 vs Group 2 : N/A : Descriptive data for each participant In both groups: AROM, PROM, MAS, BMRC improved ABC checklist improved only in Group 1	5	
Seifart et al. ⁴⁵ 2010	Randomised single subject trial	Functional electrical stimulation (FES)	Stimulation of gastrocnemius and tibialis anterior beginning at 5 different times post injection into TS for 4 weeks home programme: at 1, 7, 14, 32, 35 days. No control group	5 children Cerebral Palsy Unilateral GMFCS 1 Mean age: 4.5 years	Isometric muscle strength of the ankle plantar flexors and dorsiflexors (Hand-held dynamometer)	Self-selected walking speed (10m Walking test)		2 months	Mean change Group 1 vs Group 2 : N/A : Descriptive data for each participant Some increase in isometric plantar flexor strength No change in walking	9	
Rha et al. ⁴⁶ 2008	RCT	Electrical stimulation (ES)	Stimulation of gastrocnemius 7 consecutive days after injection and a sham stimulation on the other side 30 min a day 2 groups: HFES (25Hz) and LFES (4Hz) Biphasic rectangular current, 0.25 ms pulse	23 children Cerebral Palsy Bilateral GMFCS 1,2,3 Mean age: 46 months	Compound Motor Action Potentials for gastrocnemius Spasticity (MAS, TS)			1 month	Mean change Group 1 vs Group 2 : N/A : no control group Earlier denervating action of BTX-A Not correlated to clinical reduction of spasticity	8	
Detrembleur et al. ⁴⁷ 2001	RCT	Electrical stimulation (ES)	Stimulation in injected muscles beginning on the day of BTX 30 minutes, 6 times a day for 3 days Continuous trains of current pulses (20 Hz, 0.2 msec, 50-90 mA) Group 1 : Group 1 : BTX + ES Group 2 : BTX alone	12 children Cerebral Palsy Unilateral / Bilateral GMFCS 1,2 Median age: 5 years	Spasticity: MAS PROM Ankle muscle stiffness	Gait: Physician Rating Scale 3D-Instrumented Gait analysis		6 months	Mean change Group 1 vs Group 2 : N/A : descriptive data for group1 are not presented → adjunct ES had no significant effect (p>0.05) on clinical measurements, ankle stiffness and gait variables	8	
Williams et al. ⁴⁸ 2013	Cross-comparison design with a 6-month preintervention controlled period	Resistance training (RT)	Lower limb Home-based training programme 3 times a week for 10 weeks Progressive strengthening exercises Initially, work on motor control and then more complex movements and functional tasks Group 1 : BTX+RT Group 2 : BTX alone 2 subgroups : either pre or post BTX RT	15 children Cerebral Palsy Bilateral GMFCS 1,2 Mean age: 8 years	Spasticity: MAS Selective Control Assessment of the lower extremity Muscle isometric and isokinetic strength of the knee and ankle muscles Muscle Volume (MRI)	GAS		3 months	Many evaluations for many muscles : MAS : significantly reduced after BTX. No significant change over the strength training for either group Strength : significant isokinetic strength gains in the intervention period compared to the control period GAS : significant improvement compared to the control period MV : significant improvement in all assessed muscles compared to the control period	9	
Eivrum et al. ⁴⁹ 2012	RCT	Resistance training (RT)	Upper limb RT: 3 times a week for 8 weeks with a physiotherapist 0-40 minutes core strengthening and singlejoint resistance training with increasing intensity by 0.25 - 0.5kg Group 1 : BTX+RT Group 2 : BTX alone	10 children Cerebral Palsy Unilateral / Bilateral MACS 2 Mean age: 13.4 years	AROM of elbow and forearm Muscle tone and strength in the elbow and forearm Isometric grip force	Hand and arm use: Melbourne AHA		9 months	Mean change Group 1 vs Group 2 : MAS : 1.5 (N/A) vs 1.2 (N/A) Dorsiflexion MT : N/A no change Plantarflexion MT : +0.12 (N/A) vs -0.11 (N/A) † GMFM : +1.8 (N/A) vs +4.2 (N/A) No changes in postural control, kinematics and gait parameters in both groups	6	
Bandholm et al. ⁵⁰ 2012	A randomised pilot study	Resistance training (RT)	Lower limb RT: 2 times per week for 12 weeks 10 minutes of gait and balance 5 minutes of stretching 15 minutes of progressive resistance training Group 1 : BTX + RT Group 2 : BTX alone	14 children Cerebral Palsy Unilateral GMFCS : 1 Mean age: 9.5 years	Spasticity: MAS Dorsiflexion Maximal Torque (MT) Plantarflexion MT	Balance (biomechanical force plate) GMFM 3D - Gait analysis (Vicon)		3 months	Mean change Group 1 vs Group 2 : MAS : 1.5 (N/A) vs 1.2 (N/A) Dorsiflexion MT : N/A no change Plantarflexion MT : +0.12 (N/A) vs -0.11 (N/A) † GMFM : +1.8 (N/A) vs +4.2 (N/A) No changes in postural control, kinematics and gait parameters in both groups	9	
Speth et al. ⁵¹ 2015	Multicentre RCT	Bimanual task-oriented therapy (BTOT)	30 min of physiotherapy (PT) and one hour of occupational therapy (OT) 2 times a week, for 12 weeks Bimanual goals were set using the Canadian Occupational Performance Measure (COPM) Group 1 : BTX+BTOT Group 2 : BTX alone Group 3 : BTOT alone	35 children, Cerebral Palsy Unilateral MACS levels 1-3 Mean age: 7.1 years	Spasticity in wrist and elbow: SPAT PROM and AROM of wrist, elbow and thumb Grip strength	Functional grip strength		6 months	Pretreatment : Functional strength : One hand : 162±173 vs 95.0±118 vs 131±163 Two hands : 330±2272 vs 3210±4121 vs 3182±1604 Posttreatment : Functional strength : One hand : 215±14N/A vs N/A vs 309±1N/A Two hands : 519±2N/A vs N/A vs 530±2N/A Mean change Group 1 vs Group 3 : Functional strength : One hand : 62(N/A) vs 178(N/A)* Two hands : 1188(N/A) vs 2120(N/A)	8	
Jianjun et al. ⁵² 2013	RCT	Rehabilitation program	Group 1: BTX-A injection + 2 h/day rehabilitation Group 2 : BTX-A injection + <2 h/day rehabilitation	244 Cerebral Palsy Localisation? GMFCS? Mean age: 6.4 years	Spasticity: MAS	GMFM		1 year	Pretreatment : MAS : 2.6±1.0 vs 2.7±1.1 GMFM : 45.7±8.5 vs 44.5±9.1 Posttreatment : MAS : 1.6±0.5* vs 1.6±0.6* at 1 month post BTX GMFM : 60.9±10.6* vs 56.0±9.0* at 1 year post BTX Mean change Group 1 vs Group 2 : MAS : N/A GMFM : 15.2±3.5 vs 11.5±3.2 †	8	
Park et al. ⁵³ 2009	Case-controlled non randomised study	Modified constraint-induced movement therapy (mCIMT)	Group 1 : A combined therapy of mCIMT and BTX-A injections during 3 weeks Group 2 : BTX-A injections only	29 children Cerebral Palsy Unilateral MACS? Median age: 4 years	Spasticity: MAS, MTS PROM	Upper Limb Physician's rating scale (ULPRS) How Often scale and How Well scale in the revised Paediatric Motor Activity Log (PMAL)		3 weeks	Pretreatment : MAS : 1.5(0.3 - 2.3) vs 1.5(0.3 - 2.5) R1 : 11.3(- 37.5 - 58.8) vs 22.5(- 23.8 - 55.0) How often scale : 0.6(0.0 - 1.5) vs 0.6(0.2 - 1.2) How well scale : 0.5(0.0 - 1.4) vs 0.6(0.0 - 1.6) Posttreatment : MAS : 1.0(0.0 - 2.0) * vs 1.3(0.0 - 2.5) * R1 : 37.5(6.3 - 63.8) * vs 36.3 (0.0 - 65.0) * How often scale : 1.2(0.0 - 1.5) * vs 0.8(0.3 - 1.4) † How well scale : 1.2(0.0 - 1.9) * vs 0.5(0.0 - 1.6) Mean change Group 1 vs Group 2 : MAS : 0.4(0.0 - 1.0) vs 0.3(0.0 - 1.0) R1 : 10.6(0.0 - 52.5) vs 10.0(- 6.3 - 43.8) How often scale : 0.4(- 0.0 - 0.9) vs 0.0(- 0.1 - 0.2) † How well scale : 0.8(0.0 - 1.2) vs 0.0(- 0.3 - 0.9) †	17	

Adjunct therapy	Comments
<p>Serial Casting</p> <p>Casting</p> <p>Resistance training (RT)</p> <p>Rehabilitation programme</p> <p>BTX-A alone</p> <p>Electrical stimulation</p>	<p>Variability of protocols concerning timing of serial casting</p> <p>In lower limb no studies exist involving other joints than the ankle</p> <p>Recommended – Do it</p> <p>No argument today in favour of serial casting rather than casting</p> <p>Recommended – Do it</p> <p>Requires more studies</p> <p>Remaining issues:</p> <p>Timing of RT?</p> <p>Intensity?</p> <p>Task oriented to improve function?</p> <p>Probably recommended – Probably do it</p> <p>To date very few studies are available.</p> <p>Encouraging results concerning task-orientated reeducation but requires more studies</p> <p>Probably recommended - Probably do it</p> <p>Not recommended - Don't do it</p> <p>Not recommended - Don't do it</p>