Stability, spectra and structure of the copper(II) chloride complexes in acetic acid.

Mustayeen A. Khan, J. Meullemestre, M.J. Schwing, F. Vierling

► To cite this version:

Mustayeen A. Khan, J. Meullemestre, M.J. Schwing, F. Vierling. Stability, spectra and structure of the copper(II) chloride complexes in acetic acid.. Polyhedron, 1983, 2 (6), pp.459-463. 10.1016/S0277-5387(00)87093-X . hal-03163593

HAL Id: hal-03163593

<https://univ-angers.hal.science/hal-03163593>

Submitted on 10 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

STABILITY, SPECTRA AND STRUCTURE OF THE COPPER(II) CHLORIDE COMPLEXES IN ACETIC ACID

M. A. KHAN, J. MEULLEMEESTRE, M. J. SCHWING and F. VIERLING*

E.N.S.C.S., Laboratoire de Chimie Physique ERA No. 166 1, rue Blaise Pascal, 67008 Strasbourg, France

(Received 5 July 1982)

Abstract—The stability constants and the electronic spectra of four molecular chlorocomplexes formed in acetic acid solutions have been calculated. The spectrophotometric measurements were performed at wavelengths ranging from 260 to 500 nm and, in the near IR from 600 to 1400 nm. The matrix rank treatment of more than 1000 spectrophotometric data demonstrates a minimum of five absorbing species: the solvated copper(II) acetate and four chlorocomplexes. The overall stability constants of $\text{Cu}(\text{OAc})\text{Cl}$, CuCl_2 , LiCuCl_3 and Li_2CuCl_4 are respectively: $\log \beta_1 = 3.38$; $\log \beta_2 = 5.57$; $\log \beta_3 = 7.36$; $\log \beta_4 = 7.83$. The structure of the monochlorocuprate with the Cu-OAc bonding in the primary coordination sphere is supported by near-IR evidence. The bathochromic shift of the absorption maxima of the $d-d$ transition bands indicates structural changes of the complexes with a square planar configuration for $\text{Cu}(\text{OAc})\text{Cl}$ promoted to a flattened tetrahedron for Li_2CuCl_4 .

An exhaustive study of the copper(II) chloride complexes in aqueous solution was carried out by Schwing and Khan^{1,2} who concluded the presence of four mononuclear successive complexes, calculated their equilibrium constants and discussed briefly the structure of the CuCl_4^{2-} ion in aqueous solution. The emphasis now seems to have shifted to the study of these copper(II) chlorocomplexes in nonaqueous solutions. This is mainly because of very small values of the overall equilibrium constants, β_i , in water, in contrast to their very high values in nonaqueous solvents. Moreover in water there is an overlapping of the characteristic bands of the complexes, which are well separated in the latter solvents.

Furlani and Morpurgo³ suggest a flattened tetrahedral (D_{2d} symmetry) structure for tetrachlorocuprates after studies in solid state and such organic solvents as acetonitrile, nitromethane, ethylalcohol, dimethylformamide (DMF) and dimethylsulphoxide (Me_2SO). Manahan and Iwamoto⁴ have proposed very high values for the equilibrium constants of CuCl^+ and CuCl_2 in acetonitrile.

Gutman *et al.* have shown spectrophotometrically the existence of four chlorocomplexes in DMF⁵ in acetonitrile and in trimethylphosphate⁶ at the same time postulating a very small equilibrium constant for CuCl_2 . On the other hand Courtot *et al.*⁷ using potentiometric and spectrophotometric means propose three copper(II) chlorocomplexes, CuCl^+ , CuCl_2 , CuCl_3^- in Me_2SO . Scharff⁸ as also proposed the existence of these complexes in propylene carbonate.

Very recent work published from our laboratory confirms the presence of three complexes postulated by Gutman *et al.* in DMF but with CuCl^+ , CuCl_3^- and CuCl_4^{2-} .⁹ The absence of the dichlorocomplex in this model is due to similar solvent and ligand donor numbers, inducing dissociation of CuCl_2 and autocomplex formation of CuCl^+ and CuCl_3^- .¹⁰ The same [1, 3, 4] model, i.e. CuCl^+ , CuCl_3^- and CuCl_4^{2-} is found in

Me_2SO instead of [1-3], i.e. CuCl^+ , CuCl_2 and CuCl_3^- suggested in Ref.⁷, but the [1-4] model, i.e. CuCl^+ , CuCl_2 , CuCl_3^- , CuCl_4^{2-} is clearly established in propylene carbonate¹¹ against the unsatisfactory [1-3] proposed in the same medium by Scharff.⁸

Beside these dissociating solvents, the non-dissociating acetic acid has also been used for the study of the copper(II) chlorocomplexes. Eswein *et al.*¹² suggest an equilibrium between the D_{4h} (square planar) and D_{2d} symmetry species for the tetrachlorocomplex in acetic acid, assigning the peak at 450 nm to D_{2d} and that at 375 nm to D_{4h} configuration. They identify the absorption maximum at 670 nm as that of cupric acetate in acetic acid but it is surprising that they do not observe any isosbestic point.

Sawada *et al.* have studied in detail the equilibria of copper(II) acetate in acetic acid with perchloric acid and lithium acetate¹³ and with hydrochloric acid and copper(II) chloride.¹⁴ They suggest that the characteristic absorption maximum of copper acetate in acetic acid is at 685 nm and observe isosbestic points at 360 and 440 nm and postulate the presence of tri- and tetrachlorocomplexes. However their work is limited to the interpretation of data obtained only at two wavelengths.

This work pertains to the formation of copper(II) chlorocomplexes in acetic acid at 25°C. The number of complexes present in the solution and their equilibrium constants were determined after a spectrophotometric study in the UV and visible region, from 270 to 500 nm and in the near-IR from 600 to 1400 nm. A quantitative interpretation of the whole spectrum (UV, vis., IR) was undertaken. All data were taken into account to calculate the equilibrium constants and the variation of the extinction coefficients of the individual copper(II) chlorocomplexes. Tests for different models of complexes present in the solution were performed in accordance with the results of the matrix rank method, mentioned in our discussion.

EXPERIMENTAL

Anhydrous acetic acid, pro analysis (Merck) was used after adding calculated amount of acetic anhydride, p.a. (Merck), with the addition of three drops of 0.001 M perchloric acid, 70% p.a.

*Author to whom correspondence should be addressed.

(Merck) and letting the solution stand overnight.¹⁵ Copper solutions were prepared from a stock solution of copper(II) acetate, p.a. (Merck). Lithium chloride p.a., (Merck) was used without any purification.

The final analytical concentration of copper was 5×10^{-4} M for measurements in UV-vis. and 3.75×10^{-3} M in IR region. In these spectrum regions chloride concentrations were varied from 5×10^{-4} to 1 M, with $l = 0.5$ cm, and from 2.5×10^{-3} to 1 M, with $l = 2$ cm, respectively.

Acetic acid was the spectrophotometric reference solution. The final solutions of copper and chloride were prepared just before the measurements. The optical densities were obtained with a CARY 17D instrument with digital interfacing and a Periferic Zip 30 rapid printer and puncher. The numerical analysis was performed on a UNIVAC 1110 computer.

RESULTS AND DISCUSSION

Acetic acid is a molecular solvent, its dielectric constant being 6.2, and therefore non-ionic in character. However two series of measurements were carried out. One with lithium perchlorate to keep the molecular environment constant and equal to 1 M and the other, without lithium perchlorate. The absorption spectra of the two series in UV, visible and IR being identical, the simpler of the two equilibria, i.e. the direct mixing of copper(II) acetate and lithium chloride solutions was retained for the study.

(a) Spectra of copper(II) chloride in acetic acid

When chloride is added to the copper(II) solution in acetic acid the colour changes from blue to greenish yellow, and eventually to bright yellow with the increase in chloride concentration. The absorption variations with respect to the chloride concentration in UV, visible and IR are presented in Fig. 1.

The free solvated copper(II) acetate has an absorption maximum at about 252 nm. The addition of chloride shows the formation of a peak at 300 nm which undergoes a hypsochromic effect as the chloride concentration increases with a shift to 280 nm. This maximum is not

Gaussian suggesting a superposition of two maxima. For chloride concentrations much higher than 0.02 M there appears a maximum at 445 nm and for concentrations higher than 0.05 M another at 380 nm. These two maxima coalesce at 385 nm at still higher concentrations. Three isobestic points are observed at 302, 357 and 437 nm. Sawada *et al.*¹⁴ have reported only two isobestic points at 360 and 440 nm whereas the work of Eswein *et al.* is conspicuous by its statement that "no isobestic points are observed". It is to be remarked that this latter work was carried out by preparing chloride solutions by dissolving the chloride salt first in small quantity of water before adjusting the volume with acetic acid. In our case it was observed that the addition of small quantities of water has a profound effect on the complexation, a few drops being sufficient to render the solution colourless from greenish yellow or bright yellow. This aquation effect has also been noted by Sawada and Tanaka.¹⁶

In the IR region the characteristic maximum of copper(II) acetate in acetic acid is observed at 680 nm ($198 \text{ M}^{-1} \text{ cm}^{-1}$). This peak which has been variously reported at 650¹⁵, 670¹² and 685 nm¹⁴ disappears as the chloride concentration increases and shifts to 1060 nm. This shift of 380 nm of the solvated copper(II) ion in acetic acid, from 680 to 1060 nm, is remarkably similar to the bathochromic shift in DMF,⁹ propylene carbonate and Me_2SO .¹¹

(b) Equilibrium in solution

Copper(II) acetate in acetic acid has been reported to have a maximum at 365 nm which has been ascribed to copper-copper bond by Yamada *et al.*¹⁷ and Martin and Whitley,¹⁸ indicating that the salt is dimerised in solution. This is confirmed by Sawada¹³ who has located this maximum at 370 nm. This maximum although not visible in our work because of low copper concentration in UV-visible, appears distinctly at 375 nm with copper concentration 3.75×10^{-3} M. In view of this and the fact that acetic acid being a non-dissociating solvent, molecular

Fig. 1. Experimental spectra of copper(II) chloride solutions: spectrum/ $10^3 \times [\text{Cl}^-]_{\text{anal}}$. UV-visible: $[\text{Cu}(\text{OAc})_2]_{\text{anal}} = 5 \times 10^{-4}$ M; $l = 0.5$ cm 1/0, 2/0.5, 3/1, 4/2, 5/3, 6/5, 7/10, 8/30, 9/70, 10/300, 11/700, 12/1000, 13/1300. near-IR: $[\text{Cu}(\text{OAc})_2]_{\text{anal}} = 3.75 \times 10^{-3}$ M; $l = 2$ cm 1/0, 2/2.5, 3/5, 4/10, 5/20, 6/40, 7/200, 8/1000.

The same 1
(2) for the
to the decc
in the c
peak at 13
previously

1 C 1 (2) (3) (4) (5) C C 1 2

association occurs and the following equilibria are proposed:

The copper(II) acetate has a dimeric structure in acetic acid but the copper(II) chlorocomplexes are monomeric.¹⁴ The formulation of the monochlorocomplex as $\text{Cu}(\text{OAc})\text{Cl}$ is supported by the near-IR evidence and is discussed further on.

(c) Experimental data and numerical analysis

The optical density variations as a function of chloride concentration were plotted from 260 to 500 nm, at intervals of 10 nm. These plots were used to obtain N optical density values for each selected wavelength for computation. Eventually a set of 24 wavelengths in the UV-visible range with $N = 28$ for each wavelength were considered. In the near IR region 20 wavelengths, from 600 to 1400 nm, at intervals of 40 nm, with $N = 24$ for each wavelength, were taken into account.

A matrix rank treatment method was performed for determining the minimum number of absorbing species present in the solution¹⁹ which requires a set of $N \times L$ data, where N is the number of solutions and L the number of wavelengths, with N imperatively greater than L . The interpretation of 1152 spectrophotometric data indicated a minimum of four species: the solvated copper(II) acetate and at least three copper(II) chlorocomplexes.

(d) Calculated models and equilibrium constants

The theoretical expressions and the mathematical treatment has been explained in our earlier publication.⁹ The computation was performed at three different mononuclear complexes models: the [1-3], the [1, 3, 4] and the [1-4] model in accordance with the matrix rank analysis results. Table 1 summarises the results of the numerical analysis of the three different models:

The quadratic mean, σ , for the first two models has a value of 0.128 and 0.130 respectively. On the other hand a far better value of $\sigma = 0.054$ is obtained for the model [1-4]. Consequently the existence of four mononuclear successive complexes is proposed in acetic acid for copper(II) chloride solutions.

The stability of the complexes is much higher than in water but comparable to that found in other solvents such as DMF^9 or $\text{Me}_2\text{SO}^{11}$ despite the different associative property of these solvents this is particularly true for the overall stability constant of the monochlorocom-

plex and the stepwise formation constant of the tetrachlorocomplex. On the contrary, very higher β_1 's were calculated in the propylene carbonate solutions for an identical [1-4] model: the associative solvent effect, predicted by the very low dielectric constant of acetic acid ($\epsilon = 6$) compared to that of propylene carbonate ($\epsilon = 69$) should be one of the factors which supports these stability differences.

Incomplete formation for each of the four species, represented in Fig. 3, is found in the acetic acid system, which render impossible the β_1 determination of one separated species and explain the drastic higher value found by Sawada *et al.* for CuCl_2 ($\log \beta_2 = 15.2$).¹⁴ These authors used data at 300 nm where they do not observe any further shift of the absorption with ligand concentration increase: consequently they postulate the unique presence of $\text{Cu}(\text{OAc})\text{Cl}$ and CuCl_2 . As a matter of fact, at 300 nm the absorption maximum does not change with higher chloride concentration as seen in our experimental spectra of Fig. 1. But other absorption maxima appear, particularly at 280 and 385 nm with chloride ligand increase, which is obviously consistent with the presence of one or two other complexes. The individual electronic spectra calculation shown next in Fig. 2 indicate clearly that the extinction coefficients of LiCuCl_3 and Li_2CuCl_4 cannot be neglected at 300 nm.

On the other hand, we found stepwise formation constants K_3 and K_4 very similar (1.78 and 0.45) to those calculated by Sawada *et al.*¹⁴ (1.68 and -0.15) using data at 385 and 450 nm: at those wavelengths the absorption due to $\text{Cu}(\text{OAc})\text{Cl}$ and CuCl_2 are effectively negligible compared to those of LiCuCl_3 (Fig. 2). Another evidence for the unique presence of the latter complexes involves the presence of isobestic points at 302, 357 and 437 nm. But once again, the discussion shows the necessity to use absorption data over the whole wavelengths range for a precise determination of the best model and accurate stability constants values.

(e) Electronic spectra of the copper(II) chlorocomplexes in acetic acid

The electronic spectra of the individual chlorocomplexes for the [1-4] model in the UV, visible and the near IR are presented in Fig. 2.

Monochlorocomplex, $\text{CuCl}(\text{OAc})$. In the near IR there is a peak at 680 ($1970 \text{ M}^{-1} \text{ cm}^{-1}$) which is obviously due to the acetate ion associated with the complex. This transition is clearly characteristic of the copper-acetate bond as it is located at the same wavelength as that observed for the copper(II) acetate solution, with a comparable extinction coefficient ($680 \text{ nm} - 198 \text{ M}^{-1} \text{ cm}^{-1}$; spectrum 1—Fig. 1). Therefore the formulation of the monochlorocomplex is proposed as

Table 1. Results of the numerical analysis of the theoretical models for the acetic acid-copper(II) chloride solutions

	[1,2,3]	[1,3,4]	[1,2,3,4]
$\log \beta_1$	3.20	8.97	3.38
$\log \beta_2$	5.66	-	5.57
$\log \beta_3$	7.46	14.26	7.36
$\log \beta_4$	-	15.93	7.83
$\sigma(\text{uv-vis})$	0.128	0.130	0.054

The same plot (2) for the error to the decomposition in the case of peak at 133' previously

| Cu | (1) (2) (3) (4) (5) Co Cu | et a

Fig. 2. Calculated electronic spectra of the copper(II) chlorocomplexes in acetic acid.

$\text{CuCl}(\text{OAc})$. This complex may also present a maximum at 260 nm as in water² and DMF.⁹ Unfortunately this absorption maximum could not be ascertained because of absorption interference due to solvent below 270 nm.

It may be pointed out here that copper acetate was used instead of copper perchlorate because with the latter, at copper concentrations greater than 10^{-3} M precipitation occurs for higher chloride concentration, especially in the presence of lithium perchlorate. Had copper perchlorate been used it would have rendered difficult the exploration of the spectrum in the near IR region where, for copper concentration less than 10^{-3} M the absorption is very low.

Dichlorocomplex, CuCl_2 . The absorption maximum of this complex is centred at 300 nm ($455 \text{ M}^{-1} \text{ cm}^{-1}$) in the UV region and can be compared to that at 315 nm in propylene carbonate. In the IR, CuCl_2 has a $\lambda_{\text{max}} = 800 \text{ nm}$ ($65 \text{ M}^{-1} \text{ cm}^{-1}$).

Trichlorocomplex, CuCl_3Li . It presents three maxima,

one each in UV, visible and IR, at 300 ($3150 \text{ M}^{-1} \text{ cm}^{-1}$), 450 ($900 \text{ M}^{-1} \text{ cm}^{-1}$) and 910 nm ($85 \text{ M}^{-1} \text{ cm}^{-1}$). The peaks in UV and visible are comparable to those in DMF and Me_2SO , but differ by 20 nm from those in propylene carbonate.

Tetrachlorocomplex, CuCl_4Li_2 . The tetrachlorocomplex has an intense absorption maximum at 280 nm ($5160 \text{ M}^{-1} \text{ cm}^{-1}$) and another at 388 nm ($2170 \text{ M}^{-1} \text{ cm}^{-1}$). These two bands, with slight differences, have been widely reported in many previous works. It is probable that the differences in the position of these bands are due to different solvents used in each work. There seems to be another hidden band of feeble intensity at 490 nm ($180 \text{ M}^{-1} \text{ cm}^{-1}$). In the IR the tetrachlorocomplex peaks out at 1080 nm ($120 \text{ M}^{-1} \text{ cm}^{-1}$).

(f) Isosbestic points

A very good harmony is noted between experimental and calculated isosbestic points.

Fig. 3. Formation curves of the copper(II) chlorocomplexes in acetic acid.

The same F
(2) for the e
to the deco
In the cu
peak at 13:
previously

1 Cc | (1) (2) (3) (4) (5) Co Cu | et i

Isosbestic points (λ in nm)	
Experimental	Calculated
302	303
357	354
437	437

(g) Formation and structure

Figure 3 represents the formation curves of the four copper(II) chlorocomplexes showing a very different pattern than either in water or in propylene carbonate. It is seen that for dilute solutions, chloride concentrations as low as 0.025 M, the fourth complex begins to form in appreciable quantities.

It is proposed that the tetrachlorocomplex has a D_{2d} symmetry in acetic acid. This can be inferred by the presence of a probable charge transfer band at 490 nm and the absence of any $d-d$ transition band below 1000 nm. It is recalled that the ion of D_{4h} symmetry presents normally two bands in this region, one between 700 and 800 nm and the other between 900 and 950 nm whereas the configuration D_{2d} presents absorption maxima at 1100 nm and above.²⁰ The presence of a band at 1080 nm and the absence of any band between 700 and 950 nm are favorable arguments for a D_{2d} structure of the tetrachlorocomplex in acetic acid.

REFERENCES

- ¹M. J. Schwing-Weill, *Bull. Soc. Chim. Fr.* 1973, 3, 823.
- ²M. A. Khan and M. J. Schwing-Weill, *Inorg. Chem.* 1976, 15, 2202.
- ³C. Furlani and G. Morpurgo, *Theor. Chim. Acta* 1963, 1, 102.
- ⁴S. Manahan and R. T. Iwamoto, *Inorg. Chem.* 1965, 4(10), 1409.
- ⁵H. Hubacek, B. Stancie and V. Gutmann, *Monatsh. Chem.* 1963, 94, 1118.
- ⁶M. Baaz, V. Gutmann, G. Hampel and J. Masaguer, *Monatsh. Chem.* 1962, 93, 1416.
- ⁷A. Foll, M. Ledemez et and J. Courtot-Coupez, *J. Electroanal. Chem.* 1972, 35, 41.
- ⁸J. P. Scharff, *Bull. Soc. Chim. Fr.* 1972, 1, 43.
- ⁹M. Elleb, J. Meullemeestre, M. J. Schwing and F. Vierling, *Inorg. Chem.*, 1980, 19, 2699.
- ¹⁰V. Gutmann, *Coordination Chemistry in Non-Aqueous solvents*, p. 30. Springer-Verlag, New York (1968).
- ¹¹M. Elleb, J. Meullemeestre, M. J. Schwing and F. Vierling, *Inorg. Chem.* 1982, 21, 1477.
- ¹²R. P. Eswein, E. S. Howald, R. A. Howald and D. P. Keeton, *J. Inorg. Nucl. Chem.* 1967, 29, 437.
- ¹³K. Sawada, H. Ohtaki and M. Tanaka, *J. Inorg. Nucl. Chem.* 1972, 34, 625.
- ¹⁴K. Sawada, H. Ohtaki and M. Tanaka, *J. Inorg. Nucl. Chem.* 1972, 34, 3455.
- ¹⁵A. I. Popov, *The Chemistry of Non-Aqueous Solvents* (Edited by J. J. Lagowski, III), p. 243. Academic Press, New York (1970).
- ¹⁶K. Sawada and M. Tanaka, *J. Inorg. Nucl. Chem.* 1973, 35, 2455.
- ¹⁷S. Yamada, H. Nakamura and R. Tuschida, *Bull. Chem. Soc. Jap.* 1957, 30, 953.
- ¹⁸R. L. Martin and A. Whitley, *J. Chem. Soc.* 1958, 1394.
- ¹⁹Z. Z. Hugus and A. A. El Awady, *J. Phys. Chem.* 1971, 75, 2954.
- ²⁰M. A. Khan, Thesis. Université Louis Pasteur, Strasbourg, France, 1975.

The same pl
(2) for the er
to the decon
In the ca:
peak at 133'
previously c

| Cor | (1) (2) (3) (4) (5) CoC CuC | et al