Potentiometric study of vitamin D3 complexes with manganese(II), iron(II), iron(III) and zinc(II) in water-ethanol medium

A.L.R Mercê, B Szpoganicz, M A Khan, X Do Than, G Bouet

► To cite this version:

A.L.R Mercê, B Szpoganicz, M A Khan, X Do Than, G Bouet. Potentiometric study of vitamin D3 complexes with manganese(II), iron(II), iron(III) and zinc(II) in water-ethanol medium. *Journal of Inorganic Biochemistry*, 1999, 73 (3), pp.167-172. 10.1016/s0162-0134(99)00014-8 . hal-03199232

HAL Id: hal-03199232

<https://univ-angers.hal.science/hal-03199232>

Submitted on 16 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Potentiometric study of vitamin D₃ complexes with manganese(II), iron(II), iron(III) and zinc(II) in water–ethanol medium

A.L.R. Mercê^a, B. Szpoganicz^b, M.A. Khan^c, X. Do Thanh^d, G. Bouet^{c,*}

^a Departamento de Química, Centro Politécnico, Paraná, CP 19081, CEP 81531-990 Curitiba, PR, Brazil

^b UFSC, Departamento de Química, Campus Universitário, Trindade, CP 476, CEP 88040-900 Florianópolis, SC, Brazil

^c Chimie de Coordination, Faculté de Pharmacie, 16 Boulevard Daviers, F-49100 Angers, France

^d Physiologie-Pharmacologie, Faculté de Pharmacie, 16 Boulevard Daviers, F-49100 Angers, France

Received 20 October 1998; received in revised form 14 January 1999; accepted 15 January 1999

Abstract

Vitamin D₃ (LH) complexes with manganese(II), iron(II), iron(III) and zinc(II) were identified in water–ethanol medium (30/70). Their stability constants were determined at 298 K and at a constant ionic strength of 0.100 M using potentiometric methods. The computerisation of the experimental data showed the presence of ML (M = metal, L = deprotonated vitamin D₃) and ML₂ species in all cases; in addition, the ML₃ iron(III) complex was detected. The calculated overall stability constants β for Mn^{II}L, Fe^{II}L, Fe^{III}L and Zn^{II}L are, respectively, in logarithms, 12.4, 16.5, 28.5 and 16.5. Under the experimental conditions, the only protonated species MLH detected was with iron(III). © 1999 Elsevier Science Inc. All rights reserved.

Keywords: Potentiometry; Vitamin D₃ metallic complexes; Water–ethanol

1. Introduction

Many metallic cations are involved in human metabolism as well as in oligotherapy. For instance, transition metal ions such as manganese(II), iron(II), cobalt(II) or zinc(II) give higher enzymatic activity levels compared to magnesium(II) for some native and mutant enzymes with Mg²⁺ as natural cofactor [1]. Zinc(II), cadmium(II), cobalt(II) and magnesium(II) could readily bind to apocalciferol of *Escherichia coli* at various sites, leading to different effects upon the activity of the enzyme [2].

In a previous article [3], we determined the stability constants of vitamin D₃ (vitD) or cholecalciferol complexes involving cobalt(II), nickel(II) and copper(II) in water–ethanol medium.

In the present paper, we describe the results of the complexation of vitD with manganese(II), iron(II), iron(III) and zinc(II) in water–ethanol medium at 298 K and at a constant ionic strength (0.100 M). Usually, both potentiometric and spectrophotometric techniques could be used for computerised determinations. In the case of vitD, its insta-

bility in aqueous solution and to light exposure did not allow us to perform spectrophotometric measurements. The complexes were identified potentiometrically and their respective formation constants were calculated.

2. Experimental

2.1. Reagents

All chemicals were of analytical grade and were used without further purification. All solutions were made with a mixture of bidistilled, deionised and carbon dioxide-free water (30%, vol./vol.) and ethanol (70%) from Merck (Brazil). VitD (cholecalciferol) was purchased from Sigma-Aldrich (France). Solutions were freshly prepared prior to their use.

The solutions of zinc(II) were obtained from nitrate (Carlo-Erba, Brazil) salt and quantified by complexometric titration [4]. Those of manganese(II) chloride were made by appropriate dilutions of TitrisolTM (Merck). Anhydrous FeCl₃ (Merck) and Fe(ClO₄)₂·2H₂O (GFS Chemicals, USA) were manipulated under inert atmosphere and were weighed in the solid state. The carbonate-free KOH solutions (0.100 M) were standardised against potassium hydrogeno-

* Corresponding author. Fax: +33-241-486-733; E-mail: gilles.bouet@univ-angers.fr

phthalate. In all experiments, the ionic strength (μ) was maintained at 0.100 M by using KNO_3 (Baker and Adamson, USA) as supporting electrolyte.

The ligand solution was transferred into the reaction vessel and was titrated with the KOH solution, first in the absence of metal ion and then in its presence. The numbers of moles of vitD, Fe^{2+} and Fe^{3+} were calculated taking into account their actual weighed mass.

2.2. Method

The potentiometric titrations were carried out under inert atmosphere using 0.100 M KOH water–ethanol solutions (saturated with dinitrogen) with continuous magnetic stirring and under a purified dinitrogen flow (White-Martins, Brazil) in a 60 ml jacketed vessel at $25.0 \pm 0.1^\circ\text{C}$ (Micro-química MQBTC 99-20) with an airtight cap fitted with gas inlet and outlet tubes, glass and calomel reference electrodes (Analyser, Brazil). The $\text{p}K_w$ of water in this water–ethanol medium was determined by potentiometric titration of water–ethanol (30/70% vol./vol.) and the determined value, mean of five experiments, was 14.71 ± 0.02 [3,5]. A 20 ml Metrohm manual piston burette was used to deliver the titrant, standard CO_2 -free KOH solution, and the $\text{p}[\text{H}]$ values ($\text{p}[\text{H}] = -\log_{10}[\text{H}^+]$, a specie between square brackets denotes its molarity) were directly measured with a Micronal B-375 model pHmeter (SP, Brazil). The pHmeter was calibrated with standard adjusted ionic strength (0.100 M, KNO_3) HCl in water–ethanol (30/70% vol./vol.) solution before starting the titration. The slope of the response of the glass electrode was determined by using the data obtained from a potentiometric titration of a known volume of this standard 0.100 M in water–ethanol (30/70%, vol./vol.) [3,5,6].

VitD was kept under inert atmosphere in a Schlenke glass and was always transferred to the reaction vessel using another Schlenke glass stopped with Suba-Seal (Sigma, USA) inside a glove bag (Sigma). About 40 mg of vitD (0.1 mol) was dissolved in 10 ml absolute ethanol p.a. and then transferred via a syringe to the titration vessel, previously prepared with all other reactants and under inert atmosphere. When investigating Fe^{2+} and Fe^{3+} ions, their salts were transferred inside a glove bag from a Schlenke glass to another one stopped with Suba-Seal and, after being weighed (0.1 and 0.05 mol approximately), they were dissolved in 4 ml of absolute ethanol and taken into the reaction vessel via another syringe prior to the transfer of vitD. All volumes of ethanol used to dissolve the reactants were taken into account in the calculation of solvent percentages.

Each potentiometric titration of approximately 0.1 mmol of vitD with 0.1 and 0.05 mmol each of the studied metallic cations was carried out three times. All titrations were made after stabilisation of the first $\text{p}[\text{H}]$ value of the system which was achieved by the successive additions of 0.04 ± 0.02 ml and 0.06 ± 0.02 ml of titrant. The corresponding $\text{p}[\text{H}]$ was read and plotted to calculate the values of the overall forma-

tion constants of the systems. All titrations were carried out in the presence of HNO_3 (Merck) in 30% H_2O –70% ethanol (vol./vol.) solutions of about 0.8 M, determined by Gran's plot. This ensured that the titrations would start from $\text{p}[\text{H}]$ values under 3.0. The titration of this strong additional acid is represented by points plotted in the negative range of the x axis in Fig. 1.

2.3. Data treatment

All equilibrium constants were calculated using the micro-computer program Best 7. All mathematical details have been previously described [3,5–8].

Mole and millimole units were used to express the quantities of reagents, and the hydrolysis constants in the present solvent for all the metal ions studied were obtained from the literature [9] and were fully employed in all calculations.

The species distribution curves were drawn with the micro-computer program SPE [5–8]. In general, three titrations were made: one with the ligand alone and two others with ligand and metals in various metal-to-ligand ratios. These ratios were chosen in such a way as to favour the formation of 1:1 and 1:2 complex species (ML and ML_2 respectively) where M is the metal ion and L the OH deprotonated vitD. The solution of vitD and Fe^{3+} in a 1:3 ratio at 1×10^{-3} M could not be made because of the low solubility of this system at low $\text{p}[\text{H}]$ values. All reported results are the average of three potentiometric titration experiments.

3. Results and discussion

The average overall apparent formation constants β_n were obtained as follows from the stepwise equilibrium:

where M is the metal ion and L the deprotonated vitD, $\text{VitD}(\text{O}^-)$. In this solvent, the value of the protonation constant for the hydroxyl group in vitD is $\log K = 12.4$ [10]:

or

Fig. 1 depicts the potentiometric $\text{p}[\text{H}]$ profiles for each titration of vitD (13 points) with Mn^{2+} (12 and 15 points),

Fig. 1. Potentiometric ($p[H]$) profiles of 0.1 mmol of vitD and 0.1 and 0.05 mmol of (a) manganese(II), (b) iron(II), (c) iron(III) and (d) zinc(II).

Fe^{2+} (11 and 14 points), Fe^{3+} (50 and 57 points) and Zn^{2+} (23 and 29 points) respectively. They show that in all cases, irrespective of metal-to-ligand ratio and cation, these systems undergo rapid hydrolysis and solvolysis, preventing, in some cases, the determination of the formation constants of the complex species. Higher ratios of ligand-to-metal titrations did not prevent this immediate hydrolysis and solvolysis.

Fig. 1 shows that there is a strong depletion in the values of $p[H]$ when the ligand is in the presence of metal ions. The titration curves for Mn^{2+} complexes show an inflexion point

at $a = 3.0$ (3.0 base equivalent) and at 0.85, 3.4 and 1.4 respectively for $Fe(II)$, $Fe(III)$ and $Zn(II)$.

Table 1 summarises the logarithms of the stability constants for all the complex species detected by potentiometric titrations calculated on the basis of the equilibria (1) to (5), taking care to avoid the formation of hydroxo and/or solvo complexes. The formation constants of 1:1, 1:2 and 1:3 metal-to-ligand ratio complexes were evaluated from the $p[H]$ profiles for solutions containing 1:1 and 1:2 molar stoichiometry of cation to ligand. The species under investigation were those

Table 1

Logarithmic values ^a for the formation constants of the complexes of vitD with Mn^{2+} , Fe^{2+} , Fe^{3+} and Zn^{2+} ions at 298 K (β =overall stability constant; K =stepwise stability constant)

	Manganese(II)	Iron(II)	Iron(III)	Zinc(II)
$\log K_1 = \log \beta_1$	6.6 ± 0.1	8.5 ± 0.1	15.5 ± 0.1	8.8 ± 0.3
$\log K_H$	n.d.	n.d.	3.0 ± 0.1	n.d.
$\log K_2$	5.8 ± 0.1	8.0 ± 0.1	13.0 ± 0.1	7.7 ± 0.3
($\log \beta_2$)	(12.4)	(16.5)	(28.5)	(16.5)
$\log K_3$	n.d.	n.d.	6.2 ± 0.1	n.d.
($\log \beta_3$)			(34.7)	

^a n.d. = not detected.

expected according to common principles of coordination chemistry.

ML_3 species were only detected in the presence of the strong Lewis acid Fe^{3+} , which is able to deprotonate and hold a third molecule of vitD. Although there was formation of insoluble products at $p[H]$ values of about 6.0, there was not enough concentration of ML_3 species to be detected by potentiometric titration.

Figs. 2–5 depict the distribution diagrams for the formation constants obtained from saturated solutions with vitD and Mn^{2+} , Fe^{2+} , Fe^{3+} and Zn^{2+} respectively, with the metal ion

concentration set at 100% and calculated from $p[H] = 2$ to 12. These diagrams show the maximum formation of the complex species ML between vitD and the metallic ion for a $p[H]$ value of 9.0 (Mn^{2+}), 7.0 (Fe^{2+}), 3.0 (Fe^{3+}) and 7.0 (Zn^{2+}). The maximum formation of ML_2 complexes occurred at $p[H] = 12.0$ for Mn^{2+} , 10.0 for Fe^{2+} , 6.0 for Fe^{3+} and 11.3 for Zn^{2+} . Finally, the maximum formation of the complex ML_3 with Fe^{3+} is located at $p[H] = 10$.

The metal ions Fe^{2+} and Fe^{3+} show some complex species throughout the range of $p[H]$ under study, despite the formation of metal hydroxide precipitates at $p[H]$ values below

Fig. 2. Species distributions of solutions of vitD (L) (2.5×10^{-3} M) and Mn^{2+} (M) (2.50×10^{-3} M) for $p[H]$ values from 2.0 to 12.0. The metal concentration is set at 100%, H_{-x} indicates $[OH^-]_x$.

Fig. 3. Species distributions of saturated solutions of vitD (2.5×10^{-3} M) and Fe^{2+} (2.50×10^{-3} M) for $p[H]$ values from 2.0 to 12.0.

Fig. 4. Species distributions of saturated solutions of vitD (2.5×10^{-3} M) and Fe^{3+} (2.50×10^{-3} M) for $p[\text{H}]$ values from 2.0 to 12.0. The metal concentration is set at 100%. H_{-x} indicates $[\text{OH}^-]_x$.

Fig. 5. Species distributions of saturated solutions of vitD (2.5×10^{-3} M) and Zn^{2+} (2.50×10^{-3} M) for $p[\text{H}]$ values from 2.0 to 12.0. The metal concentration is set at 100%. H_{-x} indicates $[\text{OH}^-]_x$.

or near neutral values. The metal ions Mn^{2+} and Zn^{2+} form stable complexes above $p[\text{H}]$ values of 7.0 and 5.0, respectively.

Fe^{3+} ions led to the highest stability constants in this series (Table 1). This indicates that there is no steric hindrance in this species for which the maximum formation is centred around $p[\text{H}] = 10.2$.

Fig. 6 shows the relation between the first stability constant K_1 relative to equilibrium (1) and the atomic number of the metal used in this study and in our previous paper [3]. It may be remarked here that the tendency of the curve shows that copper(II) leads to the highest stability of the complex. This

follows the well-known Irving–Williams series [11] with the exception of the complex with Fe^{2+} . The progression of this series in alcoholic medium has been recently verified for the metal ions under consideration [12] and has been explained on the basis of the second ionisation potential of the ions. It has been shown to be valid equally in the case of dog, bovine and human serum albumin complexing copper(II) and zinc(II) [13,14]. Though our experiments were performed under inert atmosphere to prevent ferrous ions from oxidation, the high value of K_1 in the case of Fe^{2+} could be explained on the basis of an equilibrium between $\text{Fe}(\text{II})$ and $\text{Fe}(\text{III})$ ions.

Fig. 6. Logarithms of the first stability constant for some divalent cations.

Acknowledgements

The authors thank the sponsor Universidade Federal do Paraná, UFPR, Brazil.

References

- [1] J.A. Cowan, *Chem. Rev.* 98 (1998) 1067.
- [2] J.E. Coleman, K.I. Nakamura, J.F. Chlebowski, *J. Biol. Chem.* 258 (1983) 386.
- [3] A.L.R. Mercê, B. Szpoganicz, R.C. Dutra, M.A. Khan, X. Do Thanh, G. Bouet, *J. Inorg. Biochem.* 71 (1998) 87.
- [4] G. Schwarzenbach, H. Flaschka, *Complexometric Titrations*, Methuen, London, 1969, p. 256.
- [5] E.W. Schwingel, K. Arend, J. Zarling, A. Neves, B. Szpoganicz, *J. Braz. Chem. Soc.* 7 (1996) 31.
- [6] A.E. Martell, R.J. Motekaitis, *The Determination and Use of Stability Constants*, 2nd ed., VCH, New York, 1992.
- [7] Y.J. Li, A.E. Martell, *Inorg. Chim. Acta* 214 (1993) 103.
- [8] R. Delgado, Y. Sun, R.J. Motekaitis, A.E. Martell, *Inorg. Chem.* 32 (1993) 3320.
- [9] C.F. Baes Jr., R.E. Mesmer, *The Hydrolysis of Cations*, Wiley, New York, 1976.
- [10] A.E. Martell, R.M. Smith, *NIST Critical Stability Constants of Metal Complexes*, NIST Database, 46, Gaithersburg, MD, USA, 1994.
- [11] H. Irving, R.J.P. Williams, *J. Chem. Soc.* (1953) 3192.
- [12] M. Khan, G. Bouet, F. Vierling, J. Meullemeestre, M.J. Schwing, *Transition Met. Chem.* 21 (1996) 213.
- [13] J. Masuoka, J. Hegenauer, B.R. Van Dyke, P. Saltman, *J. Biol. Chem.* 268 (1993) 21553.
- [14] J. Masuoka, P. Saltman, *J. Biol. Chem.* 269 (1994) 25557.