Chronotolerance of a nickel(II) complex with the 5-methyl 2-furaldehydethiosemicarbazone ligand in male Swiss mice.

E.M. Jouad, X. Do Thanh, G. Bouet, N. Ménard, Mustayeen A. Khan

► To cite this version:

E.M. Jouad, X. Do Thanh, G. Bouet, N. Ménard, Mustayeen A. Khan. Chronotolerance of a nickel(II) complex with the 5-methyl 2-furaldehydethiosemicarbazone ligand in male Swiss mice.. *Biological Rhythm Research*, 2002, 32 (5), pp.513-520. hal-03218032

HAL Id: hal-03218032

<https://univ-angers.hal.science/hal-03218032>

Submitted on 6 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chronotolerance of a Nickel (II) Complex with the 5-Methyl 2-Furaldehyde Thiosemicarbazone Ligand in Male Swiss Mice

E.M. Jouad¹, X. Do Thanh², G. Bouet¹, N. Ménard² and M.A. Khan¹

¹Coordination chemistry, Faculty of Pharmacy, Angers Cedex, France;

²Pharmacology – Toxicology, CHU, Angers Cedex 01, France

Abstract

In vitro nickel (II) complex presents antimitotic effects. In this work, we have studied the *in vivo* seasonal effects of nickel (II), ligand and the complex $[\text{NiCl}_2(\text{M5FTSC})_2]$ in male Swiss mice. During spring, an intra peritoneal (i.p.) injection of NiCl_2 in aqueous NaCl up to $1.0 \cdot 10^{-2} \text{ mmol} \cdot \text{kg}^{-1}$ body weight (b.w.) killed 10% of the rodents after 6 days. Lethal dose 100% (LD100) was up to $1.91 \cdot 10^{-2} \text{ mmol} \cdot \text{kg}^{-1}$ b.w.; ligand was less toxic than Ni (II), while the complex was 25% lethal at $1.37 \cdot 10^{-2} \text{ mmol} \cdot \text{kg}^{-1}$ b.w. In autumn, mice were less sensitive to NiCl_2 . The ligand and the complex, on the contrary, were more toxic. This leads us to the conclusion that, *in vivo*, chronotolerance of nickel (II), ligand and complex in aqueous solution, are quite different in spring and autumn seasons.

Keywords: Male Swiss mice, nickel (II), complex, thiosemicarbazone, chronotolerance.

Introduction

Generally, *in vivo* effects studies do not take into account the rhythmic changes in internal biologic susceptibility to drugs or chemical molecules, as well as the ethology of experimental animals. Mice are rodents that are active in spring and summer and rest in autumn and winter. Many research workers such as Reinberg et al. (1991), Cambar et al. (1994), Touitou and Haus (1994) related experiments on rodents and human beings on the subject of the moment of administration of drugs and their toxic effects. Cambar et al. (1994) showed that the same dose of mercuric chloride in intra peritoneal injection (ip) induced a different mortality rate when given in November and in February. In rats, a vitamin D₃ metabolite was more active during the night than the day on renal enzymatic markers, which presented annual variations (Do

Address correspondence to: Prof. X. Do Thanh, Faculté de Pharmacie, 16 Boulevard Daviers, F-49045 Angers Cedex, France. Fax: (33) 2-41-48-67-33; E-mail: xuan.do@univ-angers.fr

et al., 1994, 1995). Effects of cobalt (II), 1,25-dihydroxyvitamin D₃ and their complex were quite different in spring and autumn (Do et al., 1996).

One of the main fields of research of our laboratory is the study of chemical and/or physiological effects of transition metals, various organic ligands and their mutual complexes. It has been reported that copper (II) is an anti-inflammatory metal and has some effects in carcinogenesis with different chronotolerances in spring and autumn (Hall et al., 1996).

We have already published the results concerning the chronotoxicity of a copper (II) complex (Do et al., 2001). In the present study, we continue our investigation on the new antimitotic metal complexes for which we have synthesised a series of complexes with nickel (II) and various furanic ligands. Preliminary *in vitro* studies revealed that these compounds presented anti-mitotic effects towards cultured colon adenocarcinoma cells (Jouad et al., 2002). It is therefore interesting to study their toxic effects in mice, in spring and autumn before further *in vivo* experiments, as knowledge of the administration time, can coincide with best effectiveness. We have chosen to perform our experiments during spring and autumn, because these seasons correspond to important hormonal variations whereas winter and summer correspond to hormonal stability. These data are useful in order to optimise both timing and dosage of any anti-mitotic molecule.

Materials and Methods

Preparation of the compounds

Nickel (II) chloride NiCl₂ · 6 H₂O (Fluka, p.a.) was used without further treatment. The ligand 5-methyl 2-furaldehyde thiosemicarbazone (M5FTSC), presented in Figure 1 and the complex [NiCl₂(M5FTSC)₂] were prepared and purified as previously described (Jouad et al., 2001). They were all dissolved in aqueous 9 g.L⁻¹ sodium chloride.

Animals

Male Swiss mice from Debré (Saint Doulchard, France) weighing from 18 to 25 g were used. They were chosen because of their regular weight gain and the absence of oestral cycle with changes in sexual hormonal status which could lead to errors in *in vivo* experiments (Laroche & Rousselet, 1990). Rodents have free access to both water and standard diet containing 0.8% Ca, 0.7% P and 3 IU.g⁻¹ vitamin D. They

Figure 1. Chemical structure of the ligand.

were placed at room temperature ($20 \pm 2^\circ\text{C}$) and exposed to light/dark cycle (12 h/12 h) alternatively. The daily cycle was inverted and the animals were kept in light during the night and in darkness during the day for 5 weeks, allowing us to observe their active phase from 8:00 a.m. to 8:00 p.m.

Experiments with nickel (II)

In addition to the control series, seven series of 10 mice were used and each rodent received a single i.p. injection ($200\ \mu\text{L} \cdot 20\ \text{g}^{-1}$ body weight, b.w.) between 10:00 and 12:00 a.m. On the 2nd of June, animals were treated as follows:

- Control series: animals received the vehicle, aqueous $9\ \text{g} \cdot \text{L}^{-1}$ NaCl.
- Series 1: $2.74 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 2: $5.48 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 3: $8.02 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 4: $12.02 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 5: $13.71 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 6: $14.77 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 7: $19.00 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$

All animals were placed again at room temperature and 12 h light/12 h dark (12L/12D) during 7 days. The same protocol was used in autumn (2nd of December).

Experiments with the ligand

Six series of 10 mice, in addition to the control series, were used and each animal received a single i.p. injection ($200\ \mu\text{L} \cdot 20\ \text{g}^{-1}$ b.w.) between 10:00 and 12:00 a.m. On the 2nd of June, rodents were treated as follows:

- Control series: animals received the vehicle, aqueous $9\ \text{g} \cdot \text{L}^{-1}$ NaCl.
- Series 1: $5.41 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 2: $10.11 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 3: $15.03 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 4: $25.14 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 5: $29.23 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$
- Series 6: $30.05 \cdot 10^{-3}\ \text{mmol} \cdot \text{kg}^{-1}$

The same protocol was used in the autumn (2nd of December).

Experiments with the complex

Seven series of 10 mice and the control series were used and each animal received a single i.p. injection ($200\ \mu\text{L} \cdot 20\ \text{g}^{-1}$ b.w.) between 10:00 and 12:00 a.m. On the 2nd of June, rodents were treated as follows:

- Control series: animals received the vehicle, aqueous 9 g.L⁻¹ NaCl.
- Series 1: 5.55.10⁻³ mmol.kg⁻¹
- Series 2: 11.00.10⁻³ mmol.kg⁻¹
- Series 3: 13.72.10⁻³ mmol.kg⁻¹
- Series 4: 16.44.10⁻³ mmol.kg⁻¹
- Series 5: 22.00.10⁻³ mmol.kg⁻¹
- Series 6: 25.22.10⁻³ mmol.kg⁻¹
- Series 7: 33.08.10⁻³ mmol.kg⁻¹

The same protocol was used in autumn (2nd of December).

Results

In spring, NaCl and nickel (II) chloride at 2.74.10⁻³ mmol.kg⁻¹ b.w. did not harm mice under study as shown in Table 1. But, 8.02.10⁻³ mmol.kg⁻¹ b.w. was LD₅₀. A dose up to 14.77.10⁻³ mmol.kg⁻¹ was the LD₁₀₀, 24 hours only after injection.

Table 1. Toxic effects of NiCl₂ in aqueous NaCl 9 g.L⁻¹ in spring.

Doses (mmol.kg ⁻¹)	Living animals 7 days after injection	
Control	10	
0.65	10	LD ₀
1.30	9	LD ₁₀
1.90	8	LD ₂₀
2.60	5	LD ₅₀
2.85	2	LD ₈₀
3.25	0	LD ₁₀₀
3.50	0	

Table 2. Toxic effects of the ligand in aqueous NaCl 9 g.L⁻¹ in spring.

Doses (mmol.kg ⁻¹)	Living animals 7 days after injection	
Control	10	
0.90	10	LD ₀
1.85	8	LD ₂₀
2.75	5	LD ₅₀
4.60	3	LD ₇₀
5.35	1	LD ₉₀
5.50	0	LD ₁₀₀
6.25	0	

LD_0 for the ligand was $5.41 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$; LD_{50} was about $15.03 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$ and LD_{100} was up to $30.05 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$

For the complex, $LD_{20} = 13.72 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$; $LD_{50} = 16.44 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$ and $LD_{100} = 33.08 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$

In autumn, the nickel solution was more effective than in spring. LD_{100} was up to $13.71 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$ The results proved that NiCl_2 was toxic in mice, more toxic in autumn than during spring (Tables 1 and 4).

The ligand, in autumn, was not tolerated, as in spring (Tables 2 and 5), while the complex in autumn killed mice more rapidly, with LD_{100} up to $25.25 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$ as compared to $33.08 \cdot 10^{-3} \text{ mmol.kg}^{-1} \text{ b.w.}$ in spring. Our experiments indicated, once more that NiCl_2 modifies its toxicity when it is complexed with the present ligand (Tables 3 and 6).

Discussion

This work shows that in male Swiss mice, tolerances of aqueous NiCl_2 were quantitatively different in spring as compared to rodents in autumn maintained under

Table 3. Toxic effects of the complex in aqueous NaCl 9 g.L^{-1} in spring.

Doses (mmol.kg^{-1})	Living animals 7 days after injection	
Control		
2.75		
5.45	10	LD_0
6.80	8	LD_{20}
8.15	5	LD_{30}
10.90	2	LD_{80}
12.50	2	
16.00	0	LD_{100}

Table 4. Toxic effects of NiCl_2 in aqueous NaCl 9 g.L^{-1} in autumn.

Doses (mmol.kg^{-1})	Living animals 7 days after injection	
Control	10	
0.65	10	LD_0
1.30	9	
1.90	9	
2.60	8	LD_{20}
2.85	4	LD_{60}
3.25	3	LD_{70}
3.50	0	LD_{100}

Table 5. Toxic effects of the ligand in aqueous NaCl 9 g.L⁻¹ in autumn.

Doses (mmol.kg ⁻¹)	Living animals 7 days after injection	
Control	10	
0.90	10	LD ₀
1.85	6	LD ₄₀
2.75	5	LD ₅₀
4.60	3	
5.35	1	LD ₉₀
5.50	0	LD ₁₀₀
6.25	0	

Table 6. Toxic effects of the complex in aqueous NaCl 9 g.L⁻¹ in autumn.

Doses (mmol.kg ⁻¹)	Living animals 7 days after injection	
Control	10	
2.75	10	LD ₀
5.45	10	LD ₀
6.80	7	LD ₃₀
8.15	5	LD ₅₀
10.90	2	LD ₈₀
12.50	0	LD ₁₀₀
16.00	0	

constant conditions. In Debré laboratories, rodents are kept under artificial light and dark (12L/12D) at 20 ± 2°C, so we maintained our animals under the same conditions, as indicated in our experimental section.

Experiments on laboratory animals during many seasons, under constant parameters such as light and temperature, showed that their seasonal rhythms are not altered. This has been reported by Cambar et al. (1994) and Reinberg et al. (1991). The discovery of clock genes such as *Per*, *Cry*, *Clock* or *B mal* enhanced the hypothesis that most biological rhythms are genetically programmed (Fukada & Ohano, 2002). According to Pevet (2000) and Novak and Zawilska (1999) seasonal rhythms could be regulated by a hormone (melatonin) secreted by the pineal gland in mammals.

Few papers showed seasonal chrono-effectiveness of drugs. This may be due to the size and the cost of such experiments (Cambar et al., 1994). Our toxic effects study, dealing with periodic changes of activity of nickel chloride and the complex could be correlated with previous works. According to Levi (1987), the tolerance of

adriamycine in rodents was maximal in autumn and less in spring as our nickel complex. Brugerolle et al. (1988) demonstrated that the acute toxicity of phenobarbital in mice varies during the year: The compound is more toxic in January and it is well supported in July.

We can explain the chronesthesia of nickel chloride and the studied complex on the basis of biovariability. All living beings present biovariability under control of a biochemical clock which could be melatonin (Pevet, 2000; Novak & Zawilska, 1999), so they could react differently to the same drugs when injected during various seasons in the same year. This has been explicitly shown in the cases of other molecules, while observing the variations of their kidney and brain tolerances (Levy, 1987; Cambar et al., 1987; Dorian et al., 1987; Do et al., 1987; Do et al., 2000).

For further experiments, we will have to take into account the importance of seasonal changes. We might hypothesise that circadian acrophases and bathyphases of the structural and functional parameters of rodents responsible for the susceptibility resistance cycle change along the year. This can be a result of an endogenous circumannual rhythm along a 12-month period, based upon melatonin influence (Pevet, 2000; Fukada & Okano, 2002).

Further studies are now in progress to test nickel (II) chloride, free ligand and the complex in rats presenting induced colorectal cancer. The above results show that their effects should be followed during different seasons of the year in order to determine the best time for optimal tolerance as well as the best period for higher therapeutic activity.

References

- Brugerolle B (1994): Chronopharmacology. In: Touitou Y, Haus E, eds., *Biological Rhythms in Clinical and Laboratory Medicine*. Berlin-Heidelberg: Springer, pp. 114–137.
- Brugerolle B, Prat M, Douille C, Dorfan P (1988): Are there circadian and circannual variations in acute toxicity of phenobarbital in mice? *Fundam Clin Pharmacol* 35: 301–304.
- Cambar J, Dorian C, Cal JC (1987): Chronobiologie et chronophysiopathologie rénale. *Path Biol* 35: 977–984.
- Cambar J, L'Aizou B, Cal JC (1994): Chronotoxicology. In: Touitou Y, Haus E, eds., *Biological Rhythms in Clinical and Laboratory Medicine*. Berlin-Heidelberg: Springer, pp. 138–150.
- Do Thanh X, Boissnard P, Khan MA, Girault A (1994): Day time and night time effects of 24,25-(OH)₂ cholecalciferol on renal alkaline phosphatase and γ -glutamyl transpeptidase activities in intact rats. *Arch IntPhysiol Biochim Biophys* 102: 221–224.
- Do Thanh X, Coquin AS, Saïd M, Khan MA, Bouet G (1996): Spring time and autumn time effects of 1,25-dihydroxy vitamin D₃, cobalt (II) and [CoCl₂(1,25-(OH)₂D₃)] complex on renal and cerebral enzymatic markers. *Biol Rhythm Res* 27: 105–112.
- Do Thanh X, Peter JC, Annaix V, Girault A, Darcy F, Breger R, Bouet G (1995): Annual variation of cerebral and renal alkaline phosphatase and γ -glutamyl transpeptidase activities in rats. *Biol Rhythm Res* 26: 486–492.

- Do Thanh X, Djebbar-Sid S, Duchêne S, Benali-Baïtich O, Bouet G, Khan MA (2001): Chronotoxicity of a copper (II) complex with a linear tetradentate ligand in mice. *Biol Rhythm Res* 32: 423–429.
- Dorian C, Catroux PH, Cambar J (1987): Mise en évidence des variations saisonnières des rythmes circadiens de la chrononéphrotoxicité de l'Amikacine chez le rat. *Path Biol* 35: 731–734.
- Fukada Y, Okano T (2002): Circadian clock system in the pineal gland. *Mol Neurobiol* 25: 19–30.
- Hall IH, Chen SY, Tajendrah KG, West DX (1996): The anti-inflammatory activity of metal complexes of heterocyclic thiosemicarbazones, 2-substituted pyridines N-oxides and 2-pyridylthioureas. *Appl Organomet Chem* 10: 485–493.
- Jouad EM, Larcher G, Allain M, Riou A, Bouet G, Khan MA, Do Thanh X (2001): Synthesis, structure and biological activity of Nickel (II) complexes of 5-methyl 2-furaldehyde thiosemicarbazone. *J Inorg Biochem* 86: 565–571.
- Jouad EM, Do Thanh X, Bouet G, Bonneau S, Khan MA (2002): *In vitro* and *in vivo* effects of $[\text{Ni}(\text{M5FTSC})_2\text{Cl}_2]$ complex in cancer: Preliminary tests. *Anticancer Research* 22: 1713–1716.
- Laroche MJ, Rousselet F (1990): La souris. In: *Les animaux de laboratoire: éthique et bonnes pratiques*. Paris: Masson, pp. 143–164.
- Levi F (1987): Chronobiologie et cancers. *Path Biol* 35: 960–968.
- Novak JZ, Zawilska JB (1999): Melatonin and its role in the circadian function. *Postepy Hig Med Dosw* 53: 445–471.
- Pevet P (2000): La mélatonine: hormone donneuse de temps. *Rhythms* 32: 11.
- Reinberg AE, Labrecque G, Smolenski MH (1991): Chronotoxicité et chronotolérance. In: Labrecque G, Somlenski MH, eds., *Chronobiologie et chronothérapie*. Paris: Flammarion Médecine-Sciences, pp. 23–24.
- Touitou Y, Haus E (1994): Biological rhythms and aging. In: Touitou Y, Haus E, eds., *Biological Rhythms in Clinical and Laboratory Medicine*. Berlin-Heidelberg: Springer, pp. 188–207.