

### Increased Oxidative Stress Induces Apoptosis in Human Cystic Fibrosis Cells

Mathilde Rottner, Simon Tual-Chalot, Hadj Mostefai, Ramaroson Andriantsitohaina, Jean-Marie Freyssinet, Maria Carmen Martinez

### ► To cite this version:

Mathilde Rottner, Simon Tual-Chalot, Hadj Mostefai, Ramaroson Andriantsitohaina, Jean-Marie Freyssinet, et al.. Increased Oxidative Stress Induces Apoptosis in Human Cystic Fibrosis Cells. PLoS ONE, 2011, 6 (9), Non spécifié. 10.1371/journal.pone.0024880. hal-03275808

### HAL Id: hal-03275808 https://univ-angers.hal.science/hal-03275808

Submitted on 1 Jul 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Increased Oxidative Stress Induces Apoptosis in Human Cystic Fibrosis Cells

# Mathilde Rottner<sup>1,2</sup>, Simon Tual-Chalot<sup>3</sup>, H. Ahmed Mostefai<sup>3</sup>, Ramaroson Andriantsitohaina<sup>3</sup>, Jean-Marie Freyssinet<sup>1,2</sup>, María Carmen Martínez<sup>3</sup>\*

1 INSERM, U770, Le Kremlin-Bicêtre, France, 2 Université de Strasbourg, Faculté de Médecine, Strasbourg, France, 3 INSERM, U694, Université d'Angers, Angers, France

#### Abstract

Oxidative stress results in deleterious cell function in pathologies associated with inflammation. Here, we investigated the generation of superoxide anion as well as the anti-oxidant defense systems related to the isoforms of superoxide dismutases (SOD) in cystic fibrosis (CF) cells. Pro-apoptotic agents induced apoptosis in CF but not in control cells that was reduced by treatment with SOD mimetic. These effects were associated with increased superoxide anion production, sensitive to the inhibition of  $I\kappa B-\alpha$  phosphorylation, in pancreatic but not tracheal CF cells, and reduced upon inhibition of either mitochondrial complex I or NADPH oxidase. CF cells exhibited reduced expression, but not activity, of both Mn-SOD and Cu/Zn-SOD when compared to control cells. Although, expression of EC-SOD was similar in normal and CF cells, its activity was reduced in CF cells, the sources being different depending on the cell type. These observations underscore a reduced anti-oxidant defense mechanism, at least in part, via diminished EC-SOD activity and regulation of Cu/Zn-SOD and Mn-SOD expressions. These data point to new therapeutic possibilities in targeting anti-oxidant pathways to reduce oxidative stress and apoptosis in CF cells.

Citation: Rottner M, Tual-Chalot S, Mostefai HA, Andriantsitohaina R, Freyssinet J-M, et al. (2011) Increased Oxidative Stress Induces Apoptosis in Human Cystic Fibrosis Cells. PLoS ONE 6(9): e24880. doi:10.1371/journal.pone.0024880

Editor: Neil A. Hotchin, University of Birmingham, United Kingdom

Received May 7, 2011; Accepted August 22, 2011; Published September 12, 2011

**Copyright:** © 2011 Rottner et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the French Association "Vaincre la Mucoviscidose" (AVL n II0532). MR was supported by a fellowship from the Fondation pour la Recherche Médicale and from GEHT doctoral fellowship, and STC by a fellowship from the Ministère de l'Enseignement Supérieur et de la Recherche (France). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

\* E-mail: carmen.martinez@univ-angers.fr

#### Introduction

Cystic fibrosis (CF) is the most prevalent inherited and lethal disease in the caucasian population. It is due to mutations in the gene encoding the cystic fibrosis transmembrane conductance regulator (CFTR) protein. Expressed in the majority of epithelial cells [1], CFTR, an ATP-dependent membrane glycoprotein, acts as a cAMP-regulated chloride channel [2], and as GSH transporter, the major anti-oxidant of the cell [3], suggesting that CFTR might regulate cellular redox status. More than 1,800 mutations have been identified resulting in a defective CFTR protein [4]. The largely prevalent mutation [*i.e.*, the deletion of residue Phe-508 ( $\Delta$ F508)] leads to a partial functional trafficking mutant that is capable of conducting chloride, but is prematurely degraded from the endoplasmic reticulum [5].

Absence of functional channel CFTR at the plasma membrane does not permit water flux, leading to dehydrated secretion in all tracts, notably airway and pancreatic tracts [6–8], and impaired secretion clearance. Obstruction of tracts results in epithelial destruction and favors the proliferation of bacteria in airways. CF is characterized by chronic inflammation even in absence of pathogens, and by the recruitment of activated neutrophils. The origin of pro-inflammatory mediator production remains obscure and appears to be the consequence of hyperactivation of NF- $\kappa$ B transcription factor and the CFTR $\Delta$ F508 retention into endoplasmic reticulum (for review see [9]). Under pathophysiological conditions, activated neutrophils and epithelial cells release highly reactive molecules towards the extracellular space, like reactive oxygen species (ROS) and reactive nitrogen species in order to attack and eliminate invasive pathogens [10,11]. However, in CF, several evidences show that the defense systems are ineffective. Indeed, mitochondrial levels of ROS are enhanced in CFTR-/- lung epithelial cell line [12] suggesting that, in CF, an increased production of ROS may be associated with cell dysfunction and the incidence of disease. Conversely, in CF airways, levels of nitric oxide (NO) have been described to be low [13] and are associated with a reduction of inducible NO synthase activity [14,15] that could favor bacterial infection [16].

In addition to excessive oxidative and nitrosative stresses, defective neutralization of ROS can also exacerbate noxious functions in CF. Superoxide anion  $(O_2^-)$  is dismutated into oxygen and hydrogen peroxide by superoxide dismutase (SOD), an endogenous cellular defense system, which decreases  $O_2^-$  levels that damage cells at excessive concentration [17]. Extracellular-SOD (EC-SOD, SOD 3), Mn-SOD (SOD 2) and Cu/Zn-SOD (SOD 1) have been described as potent inhibitors of inflammation [18,19]. Although no direct evidence has shown the involvement of deregulation of SODs in CF, the fact that EC-SOD is highly expressed in airways and up-regulated in animal models of lung injury [20], it raises the possibility that SODs, and EC-SOD in particular, may play a role in CF. We have previously reported

that CF cells displayed an exacerbated apoptosis and NF- $\kappa$ B activation, both contributing to the self-perpetuating inflammatory cycle [21]. In the present study, we investigated the involvement of oxidative stresses in the apoptotic response of CF pancreatic and tracheal cells. For this, we used two pancreatic and tracheal cell lines expressing the wild-type CFTR (PANC-1 and NT-1, respectively) or CFTR $\Delta$ F508 protein (CFPAC-1 and CFT-2, respectively). Firstly, we investigated whether oxidative stress is implicated in the exacerbated apoptotic response of CF cells using a SOD mimetic and analyzing O<sub>2</sub><sup>-</sup> production. Furthermore, we examined the origin of ROS production. Secondly, analysis of SOD expressions and activities were performed.

#### Methods

#### Reagents

Cell culture reagents, Hank's balanced salt solution (HBSS), and trypsin/EDTA were obtained from Lonza (Verviers, Belgium). Fetal calf serum (FCS) was obtained from Invitrogen (Cergy-Pontoise, France). Actinomycin D (Act D), staurosporine (St), propidium iodide (PI), type I-A RNase A, rotenone, allopurinol, apocynin, and antibody to  $\beta$ -actin were purchased from Sigma-Aldrich (St. Louis, MO). The SOD mimetic, manganese (III) tetrakis (1-methyl-4-pyridyl) porphyrin pentachloride (MnTMPyP) was obtained from Calbiochem (Nottingham, UK). Ripa lysis buffer was provided from Upstate Biotech (Hampshire, UK). Inhibitor of I $\kappa$ B- $\alpha$  phosphorylation (Bay 11-7082) was purchased from BioMol Research Labs, Inc. (Exeter, UK).

#### Cell culture and induction of apoptosis

The pancreatic cancer cell line PANC-1 expressing endogenous CFTR and CFPAC-1, presenting the CFTR $\Delta$ F508 mutation, were purchased from the American Type Culture Collection (Rockville, MD). PANC-1 is a human epithelioid pancreas carcinoma cell line and was grown in DMEM. CFPAC-1 is a human pancreatic adenocarcinoma cell line and was grown in IMDM. The tracheal cell line NT-1, derived from non-CF human fetus, and the CFT-2 cell line, homozygous for the  $\Delta$ F508 mutation, were a kind gift from Dr. M. Mergey (UMR S893 INSERM, Paris, France) and were grown in DMEM/F12 (1:1) [21]. All media were supplemented with 10% heat-inactivated FCS, 100 µg/ml streptomycin, and 100 U/ml penicillin. CFPAC-1, NT-1, and CFT-2 cells were incubated in humidified 5% CO<sub>2</sub> atmosphere at 37°C. PANC-1 cells were cultured at 37°C in a humidified atmosphere of 7.5% CO<sub>2</sub>, as recommended by the American Type Culture Collection. Cell viability was checked by Trypan blue exclusion. Cells were seeded at  $7.5 \times 10^4$  cells in T75 flasks. All experiments were carried out when the cells were 80-90% confluent. They were incubated in the presence or absence of Act D (0.5 µg/ml), St (0.33 nM), MnTMPyP (50 µM), Bay 11-7082 (Bay, 7.5  $\mu$ M), rotenone (5  $\mu$ M), apocynin (100  $\mu$ M) or allopurinol (50 µM) for 24 h. All agents were used at concentrations at which no cytotoxicity was observed, as deduced from Trypan blue exclusion.

## Superoxide anion (O2-) determination by electronic paramagnetic resonance (EPR)

After 24 h of apoptosis treatment, cell medium was replaced with deferoxamine-chelated Krebs-Hepes solution containing 1hydroxy-3-methoxycarbonyl-2,5,5-tetramethylpyrrolidin (CMH; Noxygen, Mainz, Germany) (500  $\mu$ M), deferoxamine (25  $\mu$ M), and diethyldithio carbamate (5  $\mu$ M) under constant temperature (37°C) for 30 minutes. Cells when then scrapped and frozen in plastic tubes and analyzed in a Dewar flask by EPR spectroscopy using a table-top x-band spectrometer Miniscope (MS200; Magnettech, Berlin, Germany), as previously described [22]. Values are expressed as amplitude of signal per protein concentration.

#### Determination of hypodiploid DNA

After treatments, culture medium was removed from cells growing in monolayers; adherent cells were trypsinized, detached, combined with floating cells from the original culture medium, and centrifuged. Cells were then fixed in 70% ethanol for at least 4 h at 4°C and washed once in 1 mM HBSS Ca<sup>2+</sup> before resuspension for 10 min in a solution containing type I-A RNase A (0.05 mg/ml) in HBSS containing 1 mM Ca<sup>2+</sup> at 37°C. PI was then added at a final concentration of 0.1 mg/ml, as previously described [21]. After 15 min in the dark at room temperature, samples were analyzed by flow cytometry using a FACScan flow cytometer (Becton Dickinson, San Jose, CA). Data acquisition (10,000 events in each case) and analysis were conducted using CELLQuest software (Becton Dickinson). The forward light scatter setting was E-01.

#### Western blot analysis

After incubation with apoptosis-inducing agents for 24 h, cells were scrapped in the presence of 400  $\mu$ l of Ripa buffer with 10  $\mu$ g/ml leupeptin, 10  $\mu$ g/ml pepstatin, 10  $\mu$ g/ml aprotinin, and 1 mM phenylmethanesulfonylfluoride. Samples containing 20  $\mu$ g proteins (Bio-Rad protein assay kit) were separated on 10% SDS-PAGE. Separated proteins were then blotted onto Hybond-ECL nitrocellulose membrane (Amersham Biosciences, Buckinghamshire, UK). Blots were probed with antibodies against Mn-SOD, Cu/Zn-SOD and EC-SOD (Stressgen, MI), and developed with horseradish peroxidase-conjugated secondary antibody. Bound antibodies were revealed by chemiluminescence (Pierce, Rockford, IL);  $\beta$ -actin staining was used as control. Enzyme levels were determined by densitometry analysis and were normalized with respect to  $\beta$ -actin.

#### Determination of SOD activities

Analysis of SOD activities were performed according to the manufacturer's instructions (Stressgen, MI). Briefly, cells were cultured in the absence or in the presence of Act D (0.5  $\mu$ g/ml) or St (0.33 nM), and after 24 h, cells were washed, detached with trypsin and washed with ice cold phosphate buffer saline. EC-SOD activity measurement was performed using culture supernatant and Cu/Zn-SOD and Mn-SOD using cell lysate. Also, Cu/Zn-SOD was isolated by adding ice-cold chloroform/ethanol (37.5/62.5 (v/v)) and its activity measured. Absorbance was read at 405 nm for 10 min at room temperature. Data were expressed as  $\mu$ g of protein/ $\mu$ l.

#### Statistical analysis

Data are represented as mean  $\pm$  SEM; *n* represents the number of experiments. Statistical analysis was carried out using Student's *t* test or non-parametric Mann-Whitney *U* test. Differences were considered statistically significant at a value of p < 0.05.

#### Results

## The SOD mimetic reduces the increased sensitivity to apoptogenic agents in cells with CFTR dysfunction

Pancreatic and tracheal cells were incubated in the absence and in the presence of the SOD mimetic MnTMPyP (50  $\mu$ M), 30 min before treatment with pro-apoptotic agents for 24 h. Staining with

PI revealed nuclei with hypodiploid DNA (sub-G1 peak) corresponding to apoptotic cells, measured by flow cytometry. As previously described [21], CF cells displayed exacerbated apoptosis in the presence of Act D or St (Fig. 1, A and B). Higher concentrations of apoptotic agents did not induce further increase of apoptosis, but an enhanced necrosis was observed (near of 60%). MnTMPyP treatment had no effect on basal apoptosis which was not significantly different between normal cells or cells with CFTR dysfunction (Fig. 1, A and B). Interestingly, MnTMPyP was able to decrease Act D- and St-induced apoptosis



Figure 1. The SOD mimetic MnTMPyP reduces the increased sensitivity to apoptogenic agents of cells with CFTR dysfunction. (A) At confluence, PANC-1 (n = 6, black bars) and CFPAC-1 (n = 6, open bars) and (B) NT-1 (n = 6, dark gray bars) and CFT-2 (n = 6, light gray bars) cells were treated with MnTMPyP (Mn) for 30 min before treatment with actinomycin D (Act D) or staurosporine (St) for 24 h, or without any treatment (CTL). Cells were permeabilized with 70% ethanol and hypodiploid DNA was quantified by the use of propidium iodide. \* p < 0.05, \*\*\* p < 0.001 significantly different from respective control cells; † p < 0.05 significantly different between in the absence and in the presence of Mn. doi:10.1371/journal.pone.0024880.g001

in CF cells. Indeed, MnTMPyP decreased hypodiploid DNA content by 29% and 58% in Act D- and St-treated CF pancreatic cells, respectively (Fig. 1A). Similar results were obtained in tracheal cells, MnTMPyP inhibited apoptosis by 62% and 73% in Act D- and St-treated CFT-2 cells, respectively (Fig. 1B). These results suggest that oxidative stress is involved in the induction of apoptosis in CF cells.

#### $O_2^-$ production in normal and CF cells

All types of cells exhibited an EPR feature of signals derived from  $CMH-O_2^{-}$  complex. Measurement of  $O_2^{-}$  production shows that, in pancreatic and tracheal normal cells, apoptotic treatment did not induce significant changes in  $O_2^{-}$  levels (Fig. 2. A and B). By contrast, Act D- or St-treated CF cells displayed an increase of  $O_2^-$  levels (Fig. 2, A and B). Treatment with the SOD mimetic MnTMPyP abolished the increase in O<sub>2</sub><sup>-</sup> levels evoked by Act D or St in CF cells (Fig. 2, A and B). To determine the sources of  $O_2^{-}$  production involved in the induction of apoptosis in CF cells, both pancreatic and tracheal CF cells were incubated in the presence of inhibitors of xanthine oxidase (allopurinol), NADPH oxidase (apocynin) or mitochondrial complex I (rotenone), and Act D-induced apoptosis was evaluated. Apoptosis induction was independent of xanthine oxidase in both pancreatic and tracheal cells. In contrast, rotenone reduced apoptosis in pancreatic CF cells and apocynin in tracheal CF cells (Table 1).

Because, in CF cells, NF- $\kappa$ B pathway is activated under basal as well as apoptotic conditions [21], we have investigated the effects of inhibition of I $\kappa$ B- $\alpha$  phosphorylation on O<sub>2</sub><sup>-</sup> production, using Bay 11-7082. Interestingly, when I $\kappa$ -B $\alpha$  phosphorylation was inhibited in pancreatic CF cells, basal O<sub>2</sub><sup>-</sup> production was not modified but the increase in O<sub>2</sub><sup>-</sup> generation induced by apoptotic agents was reduced (Fig. 2A). Surprisingly, in tracheal CF cells, inhibition of I $\kappa$ -B $\alpha$  phosphorylation induced a strong increase in O<sub>2</sub><sup>-</sup> production and blunted the response evoked by apoptotic treatment (Fig. 2B).

#### Expression of SOD in normal and CF cells

As shown in Fig. 3A–D, expression of both Cu/Zn-SOD and Mn-SOD was down-regulated in CF cells. On the one hand, proapoptotic stimuli significantly decreased Cu/Zn-SOD expression in normal pancreatic cells (PANC-1) but not in normal tracheal cells (NT-1). On the other hand, pro-apoptotic stimuli increased Mn-SOD expression in NT-1 cells but not in PANC-1. Of note was that pro-apoptotic stimuli had no effect on either Cu/Zn-SOD or Mn-SOD in CF cells. Concerning EC-SOD, no difference in expression was observed in both normal and CF cells (Fig. 3, E and F). In addition, pro-apoptotic treatment had no effect on EC-SOD expression.

#### SOD activity in normal and CF cells

Activities of intracellular SODs (Cu/Zn-SOD and Mn-SOD) and EC-SOD were measured in pancreatic and tracheal cells (Fig. 4). No differences were observed in Cu/Zn-SOD and Mn-SOD activities between normal and CF cells. Interestingly, activity of EC-SOD was lower in CF cells than in normal cells. Proapoptotic treatments did not modify SOD activities.

#### Discussion

It has been suggested that excessive ROS production accounts for a variety of the degenerative processes of some human diseases due to their deleterious effect to target cells [11,23,24]. The data reported here show that increased susceptibility of apoptosis of pancreatic and tracheal CF cells is associated with enhanced ROS



**Figure 2. Superoxide anion production by pancreatic and tracheal cell lines after pro-apoptotic treatments.** (A) At confluence, PANC-1 (n = 5, black bars) and CFPAC-1 (n = 5, open bars), and (B) NT-1 (n = 5, dark gray bars) and CFT-2 (n = 5, light gray bars) cells were treated with the SOD mimetic MnTMPyP (Mn) for 30 min or with the inhibitor of phosphorylation of  $l\kappa$ B- $\alpha$ , Bay-11702 (Bay) for 30 min before treatment with actinomycin D (ActD) or staurosporine (St) for 24 h, or without any treatment (CTL). Then, cells were incubated in the presence of superoxide anion spin trap and quantification of the amplitude of the superoxide anion-CMH complex signal was performed by electronic paramagnetic resonance. Values are expressed as units per protein concentration ( $\mu$ g/ml). \* p<0.05, \*\* p<0.01 significantly different from respective control cells; †† p<0.01 significantly different from in the absence of Bay. NS = not significant. doi:10.1371/journal.pone.0024880.g002

production, because SOD mimetic reduced apoptosis. Also, inhibition of NF- $\kappa$ B pathway strongly reduced O<sub>2</sub><sup>-</sup> production in CF pancreatic cells suggesting a key role of this pathway in the regulation of oxidative stress. Furthermore, expression of anti-oxidant defense enzymes, Cu/Zn-SOD and Mn-SOD, was down-regulated in CF cells whereas their activities were not affected by CFTR mutation or by apoptotic treatment. In addition, EC-SOD activity, but not its expression, was reduced in CF cells when compared to normal cells. Altogether, these results suggest that disruption of the balance between ROS generation via NF- $\kappa$ B inhibitor-sensitive pathway and anti-oxidant defense may account for the sustained apoptosis and pro-inflammatory profile observed in CF cells.

We have previously shown that both Act D and St induced apoptosis but not necrosis in CF cells as demonstrated by double staining Annexin V/propidium iodide and TUNEL assays [21]. In addition, no apoptosis was observed in control cells under the same experimental conditions. This was not due to a delayed apoptotic response, since hypodiploid DNA was measured in all types of cells after 12 h, 24 h, and 48 h of apoptosis stimulation and DNA fragmentation was higher in CF cells than in control cells [21]. In addition, new experiments performed in control cells using a higher concentration of apoptotic agents showed no increase of apoptosis, but an enhanced necrosis (near of 60%).

ROS are important mediators of apoptosis mainly in vascular and epithelial cells, which subsequently initiate a series of local chemical reactions and genetic alterations resulting in an amplification of the initial ROS-mediated tissue damage and/or cytotoxicity [25]. It is estimated that normal levels of ROS are **Table 1.** Effects of inhibition of NADPH oxidase by apocynin, xanthine oxidase by allopurinol and mitochondrial complex I by rotenone on actinomycin D-induced apoptosis in pancreatic (CFPAC-1) and tracheal (CFT-2) cystic fibrosis cells (n = 5).

	CFPAC-1	CFT-2
Actinomycin D		
Apocynin	0.4±0.6%	14±0.02%*
Allopurinol	4.1±0.4%	11±0.1%
Rotenone	12±0.5%*	1.1±0.4%

Data are expressed in percentage of inhibition of actinomycin D-induced apoptosis.

\*p<0.05. doi:10.1371/journal.pone.0024880.t001 В A PANC-1 **NT-1** 250 450 CFT-2 CFPAC-1 Cu/Zn-SOD expression Cu/Zn-SOD expression 200 350 150 250 100 150 50 50 0 0 St CTL Act D CTL Act D St D С 200 300 Mn-SOD expression Mn-SOD expression 150 200 100 100 50 0 0 St CTL Act D CTL Act D St F Е 200 300 EC-SOD expression EC-SOD expression 150 200 100 100 50 0 0 CTL Act D St CTL Act D St

efficiently detoxified by endogenous enzymatic ROS scavengers such as SOD [26]. However, under conditions associated with excessive ROS production, the rate of ROS generated can exceed the capacity of anti-oxidant defense mechanisms to scavenge ROS and prevent deleterious ROS-evoked reactions. Concerning CF, it has been shown that endogenous ROS and lipid peroxidation levels are higher in CFTR - / - lung when compared to wild-type (CFTR+/+) in basal conditions, despite a strong enzymatic antioxidant expression involving SOD, indicating a constitutive redox imbalance [27]. Also, increased oxidative stress is responsible to defective autophagy in CF cells resulting in the accumulation of misfolded mutant CFTR protein [28]. Here, we provide evidence that O<sub>2</sub><sup>-</sup> mediated exacerbated apoptosis in CF cells since SOD mimetic, MnTMPyP, was able to reduce apoptosis induced by Act D and St. It has been reported that, in hepatocytes, oxidative stress induced by O<sub>2</sub><sup>-</sup> activates caspases

**Figure 3. Expression of SOD in pancreatic and tracheal cells.** At confluence, PANC-1 (n = 5, black bars) and CFPAC-1 (n = 5, open bars), NT-1 (n = 5, dark gray bars) and CFT-2 (n = 5, light gray bars) cells were treated with actinomycin D (Act D) or staurosporine (St) for 24 h, or without any treatment (CTL). Five determinations yielding similar results were performed. A  $\beta$ -actin control was included. Western-Blot were performed for Cu/Zn-SOD (A, B), for Mn-SOD (C, D) and for EC-SOD (E, F). SOD expressions were quantified by densitometric analysis and measurements were normalized with respect to  $\beta$ -actin. Densitometry values are given as mean  $\pm$  SEM \* p<0.05 significantly different from respective control cells;  $\dagger p$ <0.05,  $\dagger \dagger p$ <0.01 significantly different between both types of cells. doi:10.1371/journal.pone.0024880.g003





**Figure 4. Activity of SODs in pancreatic and tracheal cells.** At confluence, PANC-1 (n = 5, black bars) and CFPAC-1 (n = 5, open bars), NT-1 (n = 5, dark gray bars) and CFT-2 (n = 5, light gray bars) cells were treated with actinomycin D (Act D) or staurosporine (St) for 24 h, or without any treatment (CTL). Activity of both Cu/Zn-SOD and Mn-SOD are measured in A and B, and activity of EC-SOD in C and D. Enzymatic activity is expressed in absorbance units (A) per total protein concentration ( $\mu g/\mu l$ ). † *p*<0.05, †† *p*<0.01, ††† *p*<0.001 significantly different between both types of cells. doi:10.1371/journal.pone.0024880.g004

and evokes mitochondria-mediated apoptosis through the involvement of the Bcl-2 family proteins [29]. Regarding the source of  $O_2^{-}$ , whereas mitochondria seem to play a role in pancreatic CF cells, inhibition of NADPH oxidase with apocynin reduced apoptosis in tracheal CF cells indicating the implication of NADPH oxidase. These results are in accordance with those showing that the main source of  $O_2^-$  in pancreatic and tracheal cells is mitochondria and NADPH oxidase, respectively. Thus, in pancreatic cells, mitochondrial complex I and III are involved in ROS generation leading to apoptosis induction [30]. In tracheal epithelial and smooth muscle cells, the increase of O<sub>2</sub><sup>-</sup> production via the activation of NADPH oxidase may exacerbate pulmonary inflammation [31,32]. However, we cannot exclude that other sources of  $O_2^-$  might be implicated in the mechanisms leading to apoptosis in CF cells. Indeed, inhibition of complex I with rotenone reduced Act D-induced apoptosis of pancreatic CF cells about  $\sim 12\%$  whereas the SOD mimetic reduced it  $\sim 30\%$ . Similar results were observed in tracheal CF cells. These results suggest that probably other sources of  $O_2^-$  are involved in the induction of apoptosis in CF cells.

Interestingly, (i) basal  $O_2^-$  production was similar in all cell types, (ii) apoptotic treatment did not modify  $O_2^-$  production in normal cells, (iii) in CF cells, apoptotic treatment enhanced  $O_2^-$  production, (iv) inhibition of NF- $\kappa$ B pathway reduced apoptosis-induced  $O_2^-$  production in pancreatic CF cells, and (v) in tracheal

CF cells, the NF- $\kappa$ B pathway seems to regulate basal production of  $O_2^{-}$ . We have previously shown that NF- $\kappa$ B pathway controls apoptosis and inflammation in CF cells [21]. Altogether these results indicate that NF- $\kappa$ B pathway, in part via stimulation of oxidative stress, plays an important role in mediating both apoptosis and inflammation in CF cells. Unexpectedly, inhibition of NF-KB pathway, in the absence of apoptosis inducers, elicited a strong increase on O<sub>2</sub><sup>-</sup> production only in tracheal CF cells, suggesting a beneficial role for NF-KB activation in tracheal, but not pancreatic, CF cells. It is possible that differences in the regulation of basal  $O_2^-$  production by NF- $\kappa B$  are related to the different profile of pro-inflammatroy secretome of both types of cells [21] or the regulation of NADPH oxidase activity by the NFκB pathway [32,33]. Under these conditions, apoptotic stimuli failed to further enhance O2<sup>-</sup> production, probably because the system was already exhausted upon blockade of NF-KB pathway. This also strengthens the hypothesis that an exacerbated negative control of O2<sup>-</sup> production via NF-KB pathway under normal conditions. Indeed, dual effects of NF-KB by exerting either protective or deleterious effect have been reported depending on the conditions [34].

EC-SOD is highly expressed in airways and up-regulated in animal models of lung injury [20]. These results raise the possibility that SODs may play a role in CF. This hypothesis was further assessed by looking at both expressions and activities of



Figure 5. Representative schema showing that, in CF cells, increased apoptosis and NF- $\kappa$ B activation are associated with high levels of oxidative stress. Apoptotic stimuli seem activate NF- $\kappa$ B pathway which regulate reactive oxygen species (ROS) production. Whereas in pancreatic CF cells ROS are derived from mitochondria (red lines), in tracheal CF cells ROS are produced mainly by NADPH oxidase (blue lines). In addition, in both types of CF cells, a reduced anti-oxidant defense mechanism at least in part via diminished EC-SOD activity and reduced Cu/Zn-SOD and Mn-SOD expressions lead to exacerbate oxidative stress.

doi:10.1371/journal.pone.0024880.g005

three isoforms of SODs. Expression of Cu/Zn-SOD and Mn-SOD isoforms was reduced in CF cells when compared to normal cells even though their activities were not modified. The fact that apoptotic treatments exerted differential effects on the two isoforms in the normal but not in CF cells suggests that these stimuli might not play a major role in controlling the expression of these enzymes in CF. By contrast, EC-SOD expression was not

#### References

- Ashcroft FM (1999) Cystic fibrosis transmembrane conductance regulator. In: Ion Channels and Disease, Academic, San Diego, California vol. 12. pp 211–232.
- Riordan JR, Chang XB (1992) CFTR, a channel with the structure of a transporter. Biochim Biophys Acta 1101: 221–222.
- Gao L, Kim KJ, Yankaskas JR, Forman HJ (1999) Abnormal glutathione transport in cystic fibrosis airway epithelia. Am J Physiol 277: L113–L118.
- Cystic Fibrosis Mutation Database. Available: www.genet.sickkids.on.ca/cftr/ StatisticsPage.html. Accessed 2011 March 29.
- Gomes-Alves P, Penque D (2010) Proteomics uncovering possible key players in F508del-CFTR processing and trafficking. Expert Rev Proteomics 7: 487–494.
- Kreindler JL (2010) Cystic fibrosis: exploiting its genetic basis in the hunt for new therapies. Pharmacol Ther 125: 219–229.
- Taylor CJ, Aswani N (2002) The pancreas in cystic fibrosis. Paediatr Respir Rev 3: 77–81.
- Boucher RC (2007) Cystic fibrosis: a disease of vulnerability to airway surface dehydration. Trends Mol Med 13: 231–240.
- Rottner M, Freyssinet JM, Martínez MC (2009) Mechanisms of the noxious inflammatory cycle in cystic fibrosis. Respir Res 10: 23.
- Folkerts G, Kloek J, Muijsers RB, Nijkamp FP (2001) Reactive nitrogen and oxygen species in airway inflammation. Eur J Pharmacol 429: 251–262.
- Martínez MC, Andriantsitohaina R (2009) Reactive nitrogen species: molecular mechanisms and potential significance in health and disease. Antioxid Redox Signal 11: 669–702.
- Velsor LW, Kariya C, Kachadourian R, Day BJ (2006) Mitochondrial oxidative stress in the lungs of cystic fibrosis transmembrane conductance regulator protein mutant mice. Am J Respir Cell Mol Biol 35: 579–586.
- Keen C, Olin AC, Edentoft A, Gronowitz E, Strandvik B (2007) Airway nitric oxide in patients with cystic fibrosis is associated with pancreatic function, Pseudomonas infection, and polyunsaturated fatty acids. Chest 131: 1857–1864.
- Dotsch J, Puls J, Klimek T, Rascher W (2002) Reduction of neuronal and inducible nitric oxide synthase gene expression in patients with cystic fibrosis. Eur Arch Otorhinolaryngol 259: 222–226.
- Texereau J, Marullo S, Hubert D, Coste J, Dusser DJ, et al. (2004) Nitric oxide synthase 1 as a potential modifier gene of decline in lung function in patients with cystic fibrosis. Thorax 59: 156–158.

modified but its activity was reduced in CF cells. Thus, although an insufficient expression of SODs could account for an increased level of  $O_2^{-}$ , it is most likely that the reduction in EC-SOD activity plays an essential role in the elevation on  $O_2^-$  levels. In agreement with the present results, Madarasi et al. [35] have shown that in plasma from patients with CF, SOD activities were significantly lower when compared with healthy subjects. Also, reduced activity of Cu/Zn-SOD has been described in mononuclear, polymorphonuclear and red cells in CF patients [36,37]. Although EC-SOD has not been studied in CF, it has been shown that EC-SOD overexpression attenuates endotoxin-induced acute lung injury [38] and EC-SOD knock-out mice are more sensitive to pulmonary inflammation than wild type mice [39] suggesting that EC-SOD limits injury in response to many pulmonary insults. Altogether, these results suggest a reduced anti-oxidant defense mechanism in CF cells, at least in part, via diminished EC-SOD activity.

In conclusion, we provide evidence that, in CFTR mutated cells, the link between increased apoptosis and NF- $\kappa$ B activation associated with inflammation results in high levels of oxidative stress (Fig. 5). These observations further underscore a reduced anti-oxidant defense mechanism at least in part via diminished EC-SOD activity and subtle regulation of Cu/Zn-SOD and Mn-SOD expression and activities. These data point to new therapeutic possibilities targeting anti-oxidant pathways to reduce oxidative stress and apoptosis in CF cells, and thus to limit pro-inflammatory response in this pathology.

#### **Author Contributions**

Conceived and designed the experiments: RA JMF MCM. Performed the experiments: MR STC HAM. Analyzed the data: MR STC HAM. Contributed reagents/materials/analysis tools: MCM. Wrote the paper: MR STC MCM.

- Darling KE, Evans TJ (2003) Effects of nitric oxide on Pseudomonas aeruginosa infection of epithelial cells from a human respiratory cell line derived from a patient with cystic fibrosis. Infect Immun 71: 2341–2349.
- Yasui K, Baba A (2006) Therapeutic potential of superoxide dismutase (SOD) for resolution of inflammation. Inflamm Res 55: 359–363.
- Bowler RP, Nicks M, Tran K, Tanner G, Chang LY, et al. (2004) Extracellular superoxide dismutase attenuates lipopolysaccharide-induced neutrophilic inflammation. Am J Respir Cell Mol Biol 31: 432–439.
- Joseph A, Li Y, Koo HC, Davis JM, Pollack S, et al. (2008) Superoxide dismutase attenuates hyperoxia-induced interleukin-8 induction via AP-1. Free Radic Biol Med 45: 1143–1149.
- Bowler RP, Arcaroli J, Abraham E, Patel M, Chang LY, et al. (2003) Evidence for extracellular superoxide dismutase as a mediator of hemorrhage-induced lung injury. Am J Physiol Lung Cell Mol Physiol 284: L680–L687.
- Rottner M, Kunzelmann C, Mergey M, Freyssinet JM, Martínez MC (2007) Exaggerated apoptosis and NF-kappaB activation in pancreatic and tracheal cystic fibrosis cells. FASEB J 21: 2939–2948.
- Tual-Chalot S, Guibert C, Muller B, Savincau JP, Andriantsitohaina R, et al. (2010) Circulating microparticles from pulmonary hypertensive rats induce endothelial dysfunction. Am J Resp Crit Care Med 182: 261–268.
- Hoshino T, Ókamoto M, Takei S, Sakazaki Y, Iwanaga T, et al. (2008) Redoxregulated mechanisms in asthma. Antioxid Redox Signal 10: 769–783.
- Leung PS, Chan YC (2009) Role of oxidative stress in pancreatic inflammation. Antioxid Redox Signal 11: 135–165.
- Finkel T, Holbrook NJ (2000) Oxidants, oxidative stress and the biology of ageing. Nature 408: 239–247.
- Thannickal VJ, Fanburg BL (2000) Reactive oxygen species in cell signaling. Am J Physiol Lung Cell Mol Physiol 279: L1005–L1028.
- Trudel S, Kelly M, Fritsch J, Nguyen-Khoa T, Thérond P, et al. (2009) Peroxiredoxin 5 fails to limit phospholipid peroxidation in lung from Cftrknockout mice subjected to oxidative challenge. PLoS ONE 4: e6075.
- Luciani A, Villella VR, Esposito S, Brunetti-Pierri N, Medina D, et al. (2010) Defective CFTR induces aggresome formation and lung inflammation in cystic fibrosis through ROS-mediated autophagy inhibition. Nat Cell Biol 12: 863–875.
- Madesh M, Zong WX, Hawkins BJ, Ramasamy S, Venkatachalam T, et al. (2009) Execution of superoxide-induced cell death by the proapoptotic Bcl-2related proteins Bid and Bak. Mol Cell Biol 29: 3099–3112.

- Pramanik KC, Boreddy SR, Srivastava SK (2011) Role of mitochondrial electron transport chain complexes in capsaicin mediated oxidative stress leading to apoptosis in pancreatic cancer cells. PLoS One 6: e20151.
- Talbot S, Lin JC, Lahjouji K, Roy JP, Sénécal J, et al. (2011) Cigarrette smokeinduced kinin B1 receptor promotes NADPH oxidase activity in cultured human alveolar epithelail cells. Peptides 32: 1447–1456.
- Lee CW, Lin CC, Lee IT, Lee HC, Yang CM (2011) Activation and induction of cytosolic phospholipase A2 by TNF-α mediated through Nox2, MAPKs, NFκB, and p300 in human tracheal smooth muscle cells. J Cell Physiol 226: 2103–2114.
- Kuo WW, Wang WJ, Lin CW, Pai P, Lai TY, et al. (2011) NADPH oxidasederived superoxide anion-induced apoptosis is mediated via the JNK-dependent activation of NF-κB in cardiomyocytes exposed to high glucose. J Cell Physiol;doi:10.1002/jcp.22847.
- Wullaert A, van Loo G, Heyninck K, Beyaert R (2007) Hepatic tumor necrosis factor signaling and nuclear factor-kappaB: effects on liver homeostasis and beyond. Endocr Rev 28: 365–386.
- Madarasi A, Lugassi A, Greiner E, Holics K, Biro L, et al. (2000) Antioxidant status in patients with cystic fibrosis. Ann Nutr Metab 44: 207–211.
- Percival SS, Bowser E, Wagner M (1995) Reduced copper enzyme activities in blood cells of children with cystic fibrosis. Am J Clin Nutr 62: 633–638.
- Percival SS, Kauwell GP, Bowser E, Wagner M (1999) Altered copper status in adult men with cystic fibrosis. J Am Coll Nutr 18: 614–619.
- Hassett P, Curley GF, Contreras M, Masterson C, Higgins BD, et al. (2011) Overexpression of pulmonary extracellular superoxide dismutase attenuates endotoxin-induced acute lung injury. Intensive Care Med. doi: 10.1007/s00134-011-2309-y.
- Manni ML, Tomai LP, Norris CA, Thomas LM, Kelley EE, et al. (2011) Extracellular superoxide dismutase in macrophages augments bacterial killing by promoting phagocytosis. Am J Pathol 178: 2752–2759.