



HAL
open science

New constraints on the Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K.-A. Cox, E.-J. Rohling, G. Schmidt, Ralf Schiebel, S. Bacon, D.-A. Winter, M. Bolshaw, Howard Spero

► **To cite this version:**

K.-A. Cox, E.-J. Rohling, G. Schmidt, Ralf Schiebel, S. Bacon, et al.. New constraints on the Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship. *Ocean Sci. Discuss.*, 2011, 8 (1), pp.39 - 53. 10.5194/osd-8-39-2011 . hal-03277980

HAL Id: hal-03277980

<https://univ-angers.hal.science/hal-03277980>

Submitted on 17 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

**Eastern
Mediterranean
 $\delta^{18}\text{O}:\delta\text{D}$ relationship**

K. A. Cox et al.

New constraints on the Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox¹, E. J. Rohling¹, G. A. Schmidt², R. Schiebel³, S. Bacon¹, D. A. Winter⁴,
M. Bolshaw¹, and H. J. Spero⁴

¹National Oceanography Centre, Southampton SO14 3ZH, UK

²NASA Goddard Institute for Space Studies, New York, NY 10025, USA

³Laboratoire des Bio-Indicateurs Actuels et Fossiles (BIAF), Univ. d'Angers, Angers, France

⁴Geology Department, Univ. of California, Davis, One Shields Avenue, Davis, CA 95616, USA

Received: 12 November 2010 – Accepted: 6 December 2010 – Published: 18 January 2011

Correspondence to: K. A. Cox (kac@noc.soton.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Previous work on oxygen and hydrogen isotope data from Eastern Mediterranean water samples has defined a mixing relationship in this region that is different from the world surface ocean. This prompted speculations about the hydrological processes in the Mediterranean region. We present new $\delta^{18}\text{O}$ and δD data from the Eastern Mediterranean region and the East Greenland Current system, spanning a wide salinity range. These data define $\delta^{18}\text{O}:\delta\text{D}$ relationships for both regions that are consistent with the world surface ocean $\delta^{18}\text{O}:\delta\text{D}$ relationship, despite the highly evaporative conditions that prevail in the Mediterranean region. These new geochemical data have suggested that the world surface ocean $\delta^{18}\text{O}:\delta\text{D}$ relationship holds throughout almost the entire global salinity range.

1 Introduction

The relationship between oxygen and hydrogen isotope ratios in the hydrological cycle is governed by the fractionation processes associated with evaporation and precipitation. Using data from a global survey of precipitation, continental runoff and polar meltwater, Craig (1961) defined a global meteoric water line (GMWL) as:

$$\delta\text{D} = 8\delta^{18}\text{O} + 10 \quad (1)$$

where δD and $\delta^{18}\text{O}$ are the conventional delta-notations for the ratios of D to ^1H and ^{18}O to ^{16}O , expressed as a per mil (‰) deviation from the international Vienna Standard Mean Ocean Water (VSMOW) reference standard.

Equilibrium fractionation of oxygen and hydrogen isotopes during evaporation and precipitation occurs proportionally with a factor of 8. The deuterium excess (*d*-excess), defined by $\delta\text{D} - 8\delta^{18}\text{O}$, occurs as a result of kinetic fractionation occurring in addition to equilibrium fractionation during evaporation. Kinetic fractionation, sometimes referred to as vapour-diffusion fractionation, occurs as a result of the differences between the

OSD

8, 39–53, 2011

Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



molecular diffusivities of each of the isotope water species in air (e.g., Craig, 1961). The diffusion fractionation factor (and therefore the kinetic effect) for H_2^{16}O relative to HD^{16}O is not 8 times greater than that for H_2^{16}O relative to H_2^{18}O , resulting in a relative excess of D, outside the factor of 8 relationship (e.g., Craig, 1961). Increased evaporation rates at higher atmospheric vapour pressures (high temperature or low relative humidity) will therefore result in a greater kinetic versus equilibrium fractionation and thus, higher d -excess values.

2 Oxygen and hydrogen isotopes in the surface ocean

In sea surface waters, the relationship between δD and $\delta^{18}\text{O}$ is similar to that of the meteoric water line:

$$\delta\text{D} = s\delta^{18}\text{O} + i \quad (2)$$

where s is the slope of the line (7.37 ± 0.17 based on a linear regression through world surface (upper 250 m) ocean data, Rohling, (2007); grey circles in Fig. 1), and i is the intercept. In the majority of the world surface ocean, i is much smaller than in the GMWL ($-0.72 \pm 0.97\%$, Rohling, 2007; see also Schmidt et al., 1999; data originally reported by Duplessy, 1970; Weiss et al., 1979; Aharon and Chappell, 1986; Östlund et al., 1987; Yobbi, 1992; and Delaygue et al., 2001) (grey circles in Fig. 1). This world surface ocean $\delta^{18}\text{O}:\delta\text{D}$ relationship is isotopically shifted to heavier values relative to the GMWL (in that it lies to the right of the GMWL in $\delta^{18}\text{O}:\delta\text{D}$ space; see Fig. 1). Generally, the water input to the sea surface lies on the meteoric water line, however, evaporation from surface waters preferentially extracts the lighter isotopes leaving the sea surface isotopically heavy relative to the GMWL. The higher evaporation rates at low latitudes (Gat, 1996) results in a greater difference between the GMWL and the world surface ocean $\delta^{18}\text{O}:\delta\text{D}$ relationship at low latitudes relative to higher latitudes. This therefore accounts for the slightly shallower slope of the world surface ocean $\delta^{18}\text{O}:\delta\text{D}$ relationship relative to the GMWL. Laboratory experiments have shown that

Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



at 20°C and varying humidity levels, the evaporation line has a slope of 4.34 (Cappa et al., 2003). Specifically, across the Mediterranean basin, the evaporation line ranges from 3.94 to 4.24 (Gat et al., 1996; hereafter G96).

3 Oxygen and hydrogen isotopes in the Eastern Mediterranean

In semi-enclosed basins, regional meteoric water lines exist, where the slope remains close to 8 but the intercept may be higher. For example, in the Eastern Mediterranean the intercept of the meteoric water line is reported to be close to 22 (Gat, 1996; Matthews et al., 2000; McGarry et al., 2004). The $\delta^{18}\text{O}:\delta\text{D}$ relationship of surface waters in highly evaporative regions, like the Mediterranean basin, might differ from the world surface ocean as a result of the increased kinetic fractionation during evaporation into dry continental air masses. G96 reported oxygen and hydrogen isotope data from Eastern Mediterranean surface waters that define a $\delta^{18}\text{O}:\delta\text{D}$ relationship that is considerably different from that of the world surface ocean. Their reported mixing line in $\delta^{18}\text{O}:\delta\text{D}$ space is nearly flat, whereby the δD values are almost constant over a 1‰ range of $\delta^{18}\text{O}$ and i has a value of 7.44‰ (grey triangles in Fig. 1). G96 postulate that such a relationship may have developed as a result of continuous evaporation and precipitation cycles in a warm atmosphere with low relative humidity.

Schmidt et al. (2007) used a coupled ocean-atmosphere model (GISS) to simulate the oxygen and hydrogen isotope distributions in the world ocean. This model, as any global model, does not have the resolution that would allow all of the regional aspects of the relevant hydrological processes to be resolved. The data from the GISS model do however give a sound indication of the water isotope distribution in the global ocean basins. The GISS modelled annual average $\delta^{18}\text{O}:\delta\text{D}$ relationship results are also shown in Fig. 1 (in blue). The model reproduces the world surface ocean observations well with respect to $\delta^{18}\text{O}$ and δD . However, there is a distinct discrepancy between the GISS modelled and the G96 d -excess values in the Eastern Mediterranean. Schmidt et al. (2007) originally suggested that this discrepancy might indicate that the model

Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



incorporates insufficient evaporation or too much mixing with non-Mediterranean air over the Eastern Mediterranean.

Here, we present new oxygen and hydrogen isotope data from water samples collected in the Eastern Mediterranean and at Cape Farewell, East Greenland (Fig. 2) Using these data compared to previously recorded data from the Eastern Mediterranean (G96) and the rest of the world surface ocean, we investigate the discrepancy between the observed and simulated $\delta^{18}\text{O}:\delta\text{D}$ relationship in the Eastern Mediterranean.

4 Data and methods

A total of 98 surface water samples were analyzed from the two cruises; 40 of the surface water samples were collected during the November–December 2001 RV *Meteor M51-3* cruise in the Eastern Mediterranean, and the remaining 58 were collected from across the East Greenland Current system during the August–September 2005 RRS Discovery D298 cruise at Cape Farewell (Fig. 2). Together, these sample sets cover a salinity range of 29.97 to 39.38 allowing the assessment of the $\delta^{18}\text{O}:\delta\text{D}$ relationship across a very wide oceanic salinity range. This allows us to verify internal consistencies in the $\delta^{18}\text{O}:\delta\text{D}$ relationships found, as well as relative to previously reported global data. Oxygen isotope ratios were analyzed on the samples using a dual inlet Finnigan MAT 251 light stable isotope ratio mass spectrometer following equilibration of 5 ml of sample with CO_2 for 10 h at 18.1 °C. The external precision for the oxygen isotope analyses is $\pm 0.10\text{‰}$ (2σ). Hydrogen isotope ratios were analyzed on the samples by means of Cr-based pyrolysis of 0.2 μl of sample in a Eurovector 3000 elemental analyzer feeding into a continuous-flow GV Instruments Isoprime mass spectrometer. The external precision for the hydrogen isotope analyses is $\pm 1.25\text{‰}$ (2σ). Three in-house water standards, which were previously calibrated against VSMOW2 (Vienna Mean Standard Ocean Water 2), SLAP2 (Standard Light Antarctic Precipitation 2) and GISP (Greenland Ice Sheet Precipitation) (IAEA, 2009), were analyzed in duplicate during each sample run in order to produce an offset correction regression to shift the

Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



isotope data to the VSMOW scale. All analyses were performed at the University of California, Davis and we have performed an inter-laboratory calibration between the National Oceanography Centre, Southampton and the University of California, Davis.

5 Results and discussion

5 Our data from Cape Farewell (hereafter D298) define a low salinity, high latitude mixing line with a slope and intercept of 7.59 ± 0.27 and $-0.48 \pm 0.27\text{‰}$, respectively (where uncertainty limits represent 95% confidence intervals; $r^2=0.98$), which are consistent with the rest of the world surface ocean data (black squares in Fig. 1). Our new data from the Eastern Mediterranean (hereafter M51-3) do not reproduce the unusual $\delta^{18}\text{O}:\delta\text{D}$ relationship reported by G96. Instead, we find a high-salinity mixing line with a slope and intercept of 6.99 ± 0.82 and $0.31 \pm 1.11\text{‰}$, respectively ($r^2=0.89$), which agree well with the rest of the world surface ocean data (black crosses in Fig. 1). Our new data from both low salinity and high salinity regions indicate that the previously observed world surface ocean $\delta^{18}\text{O}:\delta\text{D}$ relationship holds well throughout this salinity range. The G96 mixing line for Eastern Mediterranean surface waters therefore stands out as a distinct anomaly relative to both the new data presented here, and the previously published data from the rest of the world (Fig. 1). Direct comparison of the annual average GISS data from East Greenland and the Mediterranean with our data from these regions shows that the mixing relationships are similar, however, the modelled isotope ratios are more enriched in ^{18}O and D than is observed in both regions (Fig. 1). Therefore, the anomalous, nearly flat $\delta^{18}\text{O}:\delta\text{D}$ relationship found in the Mediterranean (G96) has not been reproduced in either our new isotope data or in the GISS model data from that region. Our new data from the Mediterranean, therefore, suggest that the GISS model is in much better agreement with waters from this region than was previously thought, and that the discrepancy may be with the G96 dataset.

10

15

20

25

Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



6 Conclusions

We conclude that, in light of the new Eastern Mediterranean data, there would seem to be no anomalous behaviour of the $\delta^{18}\text{O}:\delta\text{D}$ relationship for Mediterranean surface waters, contrary to the findings of G96. The Mediterranean surface water $\delta^{18}\text{O}:\delta\text{D}$ mixing line instead remains consistent with the world surface ocean mixing line at the higher salinity end and our understanding of isotopic processes, despite the highly evaporative nature of the region. The new Eastern Mediterranean $\delta^{18}\text{O}$ data presented here have very slightly lower values than the previously obtained values, suggesting that the Eastern Mediterranean Transient (EMT) of the late 80s to early 90s impacted the distribution of $\delta^{18}\text{O}$ in the Eastern Mediterranean basin (see appendix).

We have added 98 new $\delta^{18}\text{O}:\delta\text{D}$ data points to the original world total of 244 (Schmidt et al., 1999) and the new Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ data together with that from the East Greenland Current verifies that the world surface ocean relationship holds throughout most of the global oceanic salinity range, therefore affirming this as the global surface ocean $\delta^{18}\text{O}:\delta\text{D}$ relationship. Given the agreement of our new data across the salinity range with previous global data, as well as with the GISS model output, we contend that there may have been issues with the isotope measurements of G96.

Appendix A

We note that our new Mediterranean $\delta^{18}\text{O}$ values are consistent with $\delta^{18}\text{O}$ values of surface water samples collected in this region in 1995 (Gat et al., 2003). However, these new data are $\sim 0.3\%$ lower than those of G96 and compared to the $\delta^{18}\text{O}$ data of Pierre et al. (1986) and Pierre (1999) (collected 1986–1990), our new Mediterranean data are also $\sim 0.2\%$ lower (see Fig. 4). We suggest that the shift to slightly lower $\delta^{18}\text{O}$ values in this region may have resulted from the EMT, which occurred between 1987 and 1995 (e.g., Roether et al., 1996; Klein et al., 1999). During the EMT, deep-water

Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



intrusion of Aegean waters, previously prevented by the admixture of lower density water during outflow, caused a basin-wide redistribution of salinity, temperature and nutrient concentrations (Roether et al., 1996; Klein et al., 1999; Malanotte-Rizzoli et al., 1999). These changes persisted for at least a decade (Theocharis et al., 2002). If the EMT also caused a shift in $\delta^{18}\text{O}$ values and not in salinity then that suggests a change in the freshwater source rather than a change in the amount of freshwater addition to the basin. The major rivers feeding the Black Sea, and therefore the Aegean surface waters, have higher latitude catchment areas (for example the river Danube that flows from Germany to the Black Sea) than the rivers that flow directly into the Eastern Mediterranean (for example from Turkey or Egypt). These higher latitude source rivers contribute freshwater with lower $\delta^{18}\text{O}$ values (e.g., Rank et al., 1999) to the Black Sea and the exchange of surface waters between the Black Sea and the Aegean Sea then results in a surface freshwater admixture with a relatively low $\delta^{18}\text{O}$ signature in the Aegean. Additionally, the M51-3 data points in Fig. 4 from a greater depth (100 m) indicate that these waters in the Eastern Mediterranean basin have the same salinity: $\delta^{18}\text{O}$ character as the surface waters. This is not true of the values portrayed by Pierre et al. (1986) and Pierre (1999) and G96, whereby the deeper water samples define a different salinity: $\delta^{18}\text{O}$ relationship relative to the surface samples (Fig. 4), supporting our hypothesis that the EMT has influenced $\delta^{18}\text{O}$ distribution in the Mediterranean Sea. Therefore, during the EMT, when these waters penetrated and filled the deep Mediterranean, there was, not only a redistribution of salinity, but also waters with a potentially lower $\delta^{18}\text{O}$ signature were introduced into the Eastern Mediterranean Basin.

Acknowledgements. We would like to thank T. Coplen (USGS) for his advice to M. Bolshaw with regard to calibration methods. We also thank the crews and science parties of RRS *Discovery* cruise D298 and RV *Meteor* cruise 51-3 for their assistance and support during sampling. This project was supported by the UK Natural Environment Research Council's RAPID climate change programme and the German Research Foundation (Deutsche Forschungsgemeinschaft). The data presented in this paper will be submitted to the GISS oxygen isotope database (Schmidt et al., 1999).

**Eastern
Mediterranean
 $\delta^{18}\text{O}$: δD relationship**K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Aharon, P. and Chappell, J.: Oxygen isotopes, sea-level changes and the temperature history of a coral-reef environment in New Guinea over the last 105 years, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 56, 337–379, 1986.
- 5 Cappa, C. D., Hendricks, M. B., DePaolo, D. J., and Cohen, R. C.: Isotopic fractionation of water during evaporation, *J. Geophys. Res.*, 108(D16), 4525, 2003.
- Craig, H.: Isotopic variations in meteoric waters, *Science*, 133, 1702–1703, 1961.
- Delaygue, G., Bard, E., Rollion, C., Jouzel, J., Stievenard, M., Duplessy, J.-C., and Ganssen, G.: Oxygen isotope/salinity relationship in the Northern Indian Ocean, *J. Geophys. Res.*, 106, 4565–4574, 2001.
- 10 Duplessy, J.-C.: Note preliminaire sur les variations de la composition isotopiques des eaux superficielles de l'Océan Indien: la relation ^{18}O -salinité, *C. R. Acad. Sci. Paris*, 271, 1075–1078, 1970.
- Gat, J. R.: Oxygen and hydrogen isotopes in the hydrologic cycle, *Annu. Rev. Earth Pl. Sc.*, 24, 225–262, 1996.
- 15 Gat, J. R., Shemesh, A., Tziperman, E., Hecht, A., Georgopoulos, D., and Basturk, O.: The stable isotope composition of waters of the Eastern Mediterranean Sea, *J. Geophys. Res.*, 101, 6441–6451, 1996.
- Gat, J. R., Klein, B., Kushnir, Y., Roether, W., Wernli, H., Yam, R., and Shemesh, A.: Isotope composition of air moisture over the Mediterranean Sea: an index of the air-sea interaction pattern, *Tellus B*, 55(5), 953–965, 2003.
- 20 IAEA: Reference Sheet for International Measurement Standards VSMOW2, SLAP2, International Atomic Energy Agency, 2009.
- Klein, B., Roether, W., Manca, B. B., Bregant, D., Beitzel, V., Kovacevic, V., and Luchetta, A.: The large deep water transient in the Eastern Mediterranean, *Deep-Sea Res. Pt. I*, 46(3), 371–414, 1999.
- 25 Malanotte-Rizzoli, P., Manca, B. B., d'Alcala, M. R., Theocharis, A., Brenner, S., Budillon, G., and Özsoy, E.: The Eastern Mediterranean in the 80s and in the 90s: the big transition in the intermediate and deep circulations, *Dynam. Atmos. Oceans*, 29, 365–395. 1999.
- 30 Matthews, A., Ayalon, A., and Bar-Matthews, M.: D/H ratios of fluid inclusions of Soreq cave (Israel) speleothems as a guide to the Eastern Mediterranean Meteoric Line relationships in the last 120 ky, *Chem. Geol.*, 166, 183–191, 2000.

Eastern Mediterranean $\delta^{18}\text{O}$: δD relationship

K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- McGarry, S., Bar-Matthews, M., Matthews, A., Vaks, A., Schilman, B., and Ayalon, A.: Constraints on hydrological and paleotemperature variations in the Eastern Mediterranean region in the last 140 ka given by the δD values of speleothem fluid inclusions, *Quaternary Sci. Rev.*, 23, 919–934, 2004.
- 5 Östlund, H. G., Craig, H., Broecker, W. S., and Spenser, D.: *GEOSECS Atlantic, Pacific and Indian Ocean Expeditions: Shore Based Data and Graphics*, vol. 7, National Science Foundation, Washington, DC, 1987.
- Pierre, C.: The oxygen and carbon isotope distribution in the Mediterranean water masses, *Mar. Geol.*, 153(1–4), 41–55, 1999.
- 10 Pierre, C., Vergnaud-Grazzini, C., Thouron, D., and Saliège, J.-F.: Compositions isotopiques de l'oxygène et du carbone des masses d'eau en Méditerranée, *Mem. Soc. Geol. It.*, 36, 165–174, 1986.
- Rank, D., Özsoy, E., and Salihoglu, I.: Oxygen-18, deuterium and tritium in the Black Sea and the Sea of Marmara, *J. Environ. Radioactiv.*, 43(2), 231–245, 1999.
- 15 Roether, W., Manca, B. B., Klein, B., Bregant, D., Georgopoulos, D., Beitzel, V., Kovacevic, V., and Luchetta, A.: Recent changes in Eastern Mediterranean deep waters, *Science*, 271(5247), 333–335, 1996.
- Rohling, E. J.: Progress in paleosalinity: overview and presentation of a new approach, *Paleoceanography*, 22(3), PA3215, 2007.
- 20 Schmidt, G. A., Bigg, G. R., and Rohling, E. J.: Global seawater oxygen-18 database, <http://data.giss.nasa.gov/o18data/>, last access: November 2010, 1999.
- Schmidt, G. A., LeGrande, A. N., and Hoffmann, G.: Water isotope expressions of intrinsic and forced variability in a coupled ocean-atmosphere model, *J. Geophys. Res.*, 112(D10), D10103, 2007.
- 25 Theocharis, A., Klein, B., Nittis, K., and Roether, W.: Evolution and status of the Eastern Mediterranean Transient (1997–1999), *J. Marine Syst.*, 33, 91–116, 2002.
- Weiss, R. F., Östlund, H. G., and Craig, H.: Geochemical studies of the Weddell Sea, *Deep-Sea Res.*, 26, 1093–1120, 1979.
- Yobbi, D. K.: Effects of tidal stage and ground-water levels on the discharge and water quality of springs in coastal Citrus and Herando counties, Florida, *Water Resources Investigations Report 92-4096*, US Geol. Surv., Tallahassee, Florida, 1992.
- 30




Fig. 1. The $\delta^{18}\text{O}:\delta\text{D}$ mixing relationships, from top to bottom in the legend: the modern world surface (upper 250 m) ocean (WSO) (from Schmidt et al., 1999, originally reported by Duplessy, 1970; Weiss et al., 1979; Aharon and Chappell, 1986; Östlund et al., 1987; Yobbi, 1992; and Delaygue et al., 2001), the G96 Mediterranean dataset, the M51-3 Mediterranean dataset, the D298 Cape Farewell dataset, the complete GISS model data mixing relationship (Schmidt et al., 2007), the GISS model results from the Mediterranean (GISS-M) (Schmidt et al., 2007) and the GISS model results from the East Greenland region (GISS-G) (Schmidt et al., 2007). The GMWL and an example evaporation line are also plotted.

**Eastern
Mediterranean
 $\delta^{18}\text{O}:\delta\text{D}$ relationship**

K. A. Cox et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Maps of the oxygen and hydrogen isotope data station locations for RV Meteor cruise M51-3, Pierre et al. (1986), Pierre (1999) and G96 (a) and the RRS Discovery cruise D298 (b).

Eastern Mediterranean $\delta^{18}\text{O}:\delta\text{D}$ relationship

K. A. Cox et al.




Fig. 3. Salinity, $\delta^{18}\text{O}$ and δD and distribution maps of the M51-3 data (a, b, and c) and the G96 data (d, e and f).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. Salinity: $\delta^{18}\text{O}$ plot of M51-3 data (crosses and squares) compared with Pierre et al. (1986), Pierre (1999) (circles) and G96 (triangles). Grey data points denote surface water samples and black data points denote deep water samples.

**Eastern
Mediterranean
 $\delta^{18}\text{O}:\delta\text{D}$ relationship**

K. A. Cox et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

